

## Double bonanza at the LIGO gravitational wave detectors

C. S. Unnikrishnan

*Soon after the two advanced LIGO detectors were ready in September 2015 for a calibrated observation run, a giant gravitational wave burst hit the detectors and were duly recorded as nearly identical chirped signal wave spiralling up in frequency and amplitude from 35 to 250 Hz, with a relative delay of 6.9 ms, well inside the light travel distance of 10 ms between the two detectors. The peak strain amplitude touched 100 times the base sensitivity. Analysis of the 200 ms short event led to the robust conclusion that LIGO detectors saw gravitational waves from two orbiting and merging black holes of nearly 30 solar mass each, at a distance of about 1.3 billion light years. We sketch this double discovery of great scientific and astronomical importance – the terrestrial detection of gravitational waves, ushering in the era of gravitational wave astronomy, and the discovery of a stellar mass binary black hole system, observed during their orbital evolution towards merger and then to the formation of a single stable black hole, all invisible to any other form of astronomy.*

### The discoveries

The direct laboratory detection of gravitational waves (GW) was a much awaited event by the entire physics community and the expectation ranged from ‘very soon’ to ‘sometime before 2018’, after the advanced Laser Interferometer Gravitational Wave Observatory (aLIGO) detectors came into operation at their improved sensitivity and bandwidth last year. The aLIGO optical interferometer detectors are expected to reach their projected full sensitivity only in 2017, but the confirmed stability and improved range of observations during the commissioning last year led to an engineering run in August–September followed by a science run during September 2015–January 2016. These detectors essentially measure the change in the distance between two suspended mirror masses separated by  $L = 4$  km employing optical interferometry and are designed to measure a displacement  $\delta L$  a billion times smaller than the atomic size, below  $10^{-19}$  m. The gravitational wave signal is the motion of the mirrors in response to a periodic tidal field and the relative displacement  $\delta L \propto L$ . The improvement in sensitivity from the previous version ranges from a factor of 10 to 3, for gravitational waves with frequency between 30 and 1000 Hz, which corresponds to the end stage orbital frequencies of binary neutron stars and black holes. After the calibrations and tests were completed loud and clear chirped strain signals were detected, on 14 September 2015, in both the interferometer detectors with a rela-

tive delay of 6.9 ms, well within the light travel time of 10 ms between the detectors (Figure 1).


The event is designated GW150914. Tiny by even interferometric standards, the signals exceeded signal-to-noise ratio (SNR) of 23 for the aLIGO detectors, reaching a peak strain of  $10^{-21}$ , which is more than 100 times their base sensitivity. Though sophisticated waveform matching data analysis tools are developed and used in aLIGO detectors, this signal was notified within 3 min of their arrival by a simpler burst detection algorithm. Observation of similar signals in both detectors, separated by 3000 km, was already a promising indication of a genuine common source. The statistical probability for such an event occurring due to chance coincidences from noise, estimated from the rest of the data, is less than  $2 \times 10^{-7}$ . A closer look soon confirmed two amazing discoveries<sup>1</sup>, barely a week after the detectors were in their certified stable operation; one is of course that the detected signals were indeed of gravitational waves from a binary stellar system, and the other, unexpected by most, was that the source was a binary black hole system of nearly equal mass black holes of about 29 and 36 solar masses, at an estimated distance of about 410 Mpc (redshift 0.09). Thus, this loud event just happened to be exclusively for the eyes of aLIGO, unobservable in any electromagnetic spectrum. That this happened exactly hundred years after the critical months in which Einstein was completing his theory of gravitation – the general theory of

relativity – was no strange coincidence; the advanced LIGO detectors were commissioned on schedule in the centenary year with enough sensitivity for such events for the first time.


### The detectors

Gravitational waves have a quadrupolar nature with the characteristic that head on waves force free masses into motion such that masses azimuthally separated by 90 degrees on a circle move out of phase<sup>2</sup>. This makes the Michelson interferometer that measures the path difference  $\Delta L$  between two perpendicular arms a natural detector of gravitational waves (Figure 2).

The advanced LIGO detectors<sup>3</sup> are Michelson interferometers of arm length  $L$  of 4 km each, with light path folding by about 300 multiple reflections on two mirrors implemented by Fabry–Perot cavities inside the Michelson arms (Figure 3). Thus the effective length exceeds 1000 km (about a fourth of the wavelength of gravitational waves) and the minimum detectable strain is determined by the random noise of the number of photons in the input laser beam in the frequency range 50–1000 Hz and also by thermally generated noise on the suspended optical elements at lower frequencies. The advanced LIGO detectors are the result of a major upgrade from the LIGO detectors that were operated until 2010. A comparable upgrade, with emphasis on achieving similar sensitivity in an overlapping bandwidth is underway



**Figure 1.** The strain data time series from the two detectors. Hanford detector is rotated approximately 90 degrees to Livingston detector causing 180 phase shift for the quadrupolar waves.




**Figure 2.** *a*, The quadrupolar force pattern in the gravitational wave plane, which reverses signs every half-period. *b*, The simplest gravitational wave detector monitors distance between two suspended mirror masses using optical interferometry. *c*, Michelson interferometer as a natural GW detector for the quadrupolar pattern of displacements.

at the Virgo detector in Europe as well. The major improvement is in the low frequency sensitivity, by a factor 1000 at 30 Hz to about factor 3 at 100 Hz, achieved by the enormous suppression of low frequency seismic and environmental noise in the range 10–100 Hz, employing active vibration isolation and improved passive isolation as well as reduction of thermal noise with low loss fused silica suspensions and larger mirrors. Also, heavier mirrors reduced the radiation pressure noise<sup>4</sup>. The present strain sensitivity (or more precisely, the

noise limit) at 40 and 1000 Hz is about  $2 \times 10^{-23}/(\text{Hz})^{1/2}$  (in 1 Hz bandwidth) and the best sensitivity at about 200 Hz is  $8 \times 10^{-24}/(\text{Hz})^{1/2}$ , achieved with input laser power of a modest 20 W, which is enhanced at the interferometer input to 700 W by coherently reflecting back the light repeatedly (power recycling)<sup>2,3</sup>. This corresponds to relative mirror motion of  $3 \times 10^{-20} \text{ m}/(\text{Hz})^{1/2}$ . The full projected sensitivity is another factor of 3 better, to be achieved in scheduled improvements (higher laser power, for example) during this year. The process of

commissioning towards full sensitivity is a laborious and technically complicated process because already the optical power held by the two mirrors of the Fabry–Perot cavity is 100 kW and much larger power can distort the best of mirrors technically possible to fabricate, leading to unstable operation. Isolation from seismic and environmental noise is achieved by a combination of passive isolation consisting of low frequency springs and pendulums and active feedback isolation consisting of large number of sensors and actuators. The detectors at



**Figure 3.** The optical schematic of the aLIGO detectors H1 and L1. Right insert shows the strain sensitivity. From ref. 1.

Hanford, Washington and Livingston, Louisiana are separated by about 3000 km (10.1 ms light travel time) and their orientations are such that one is rotated almost 90 degrees relative to the other (Figure 3), making the quadrupolar waves generating nearly out of phase (180 degree) response<sup>1</sup>. These exquisite and elaborate GW detectors are the result of 30 years of research and development, starting with the original experiments and proposals in the 70s. The large scale interferometers with sufficient sensitivity over a very large bandwidth like the LIGO detectors evolved from a technically and conceptually complete note by R. Weiss at MIT in 1972 (ref. 5). The interferometer detector conceived and built by Moss *et al.*<sup>6</sup> in 1971 as a laboratory prototype and the refined operational version by Forward<sup>7</sup> in 1978 were already a match to Joseph Weber's pioneering and provocative resonant bar detectors and could reach displacement sensitivity below  $10^{-15}$  m/(Hz)<sup>1/2</sup>. Formal proposals for the LIGO detectors<sup>8</sup> in the

US and the Virgo detector<sup>9</sup> in Europe were initiated in 1989. aLIGO is funded by the National Science Foundation and Virgo by CNRS, France and INFN, Italy.

### Signal, waveform and noise

The source of the transverse gravitational waves, in relativistic gravitational theories and in particular in general relativity, is time dependent matter-energy quadrupole moment and the measurable strain at distance  $R$  from the source orbiting at speed  $v$  is known by the simple formula  $h = \delta L / L = GMv^2/Rc^4$ . As the system radiates waves, the orbit shrinks due to energy loss and the orbital frequency and the strain amplitude increase. Thus the expected response in a wide-band interferometer detector is a waveform with increasing amplitude and frequency, at double the orbital frequency, called a chirp. For example, one can expect a waveform ramping up in

frequency and amplitude from  $h \sim 10^{-23}$  at 50 Hz to  $h \sim 10^{-22}$  at 500 Hz for a binary neutron star inspiral and merger. The magnitude of the strain is tiny, about  $10^{-22}$  for a 10 solar mass object at 100 Mpc, at end stage orbital size of 10 Schwarzschild radius or so. Hence, the base sensitivity has to be better than  $10^{-23}$  for detecting extragalactic binary neutron stars inspiral and merger, with SNR good enough for GW astronomy. This goal is constrained by various noise sources, all of which have been tamed fairly well in the aLIGO detectors, with some margin still ahead<sup>10</sup>. While thermal and radiation pressure noise dominate at low frequency, the detector is photon shot noise limited at frequencies above 100 Hz or so. Hence, the margin available in the laser power (<180 W) is expected to improve the high frequency sensitivity by a factor 2 to 3.

Though by chance, the first detection in the aLIGO detectors happened to be much louder than the base sensitivity of the well calibrated and stable detectors

and this high SNR makes this particular event a detector and instrumentation triumph, needing very little of sophisticated and specialized data analysis for the deduction of the essential aspects about the nature and source of the gravitational waves. It was as if one hears a mild explosion while trying to listen for an imperceptible murmur. In fact, with only the fundamental formulae for the radiated energy and strain, known from standard general relativity (GR) from early days, and the evolution of the bandpass-filtered waveform one can deduce a surprisingly large amount of information already. Matching with the template waveforms calculated using relativistic formalism with corrections to Newtonian orbits and numerical relativity, along with sophisticated data analysis tools, are of course needed, and used, for the precise estimation of the parameters of the binary source.

### Checks and confirmation

LIGO detectors are designed to detect rare events with signal strength typically too small to be seen without sophisticated data analysis methods and pipelines. Hundreds of sensors monitor the environment to map out noise from electromagnetic, seismic and other environmental disturbances. There was also the rare practice of injecting a synthesized signal, by a small authorized group of LIGO scientists, that mimics a genuine GW chirp signal into the interferometer, sometimes right at the interferometer mirrors by shaking them with actuators, without such information revealed to the other members involved in the analysis of detector state and data. The goal is to test the efficacy and readiness of all the detection and analysis pipelines and software. This is called a blind injection and the parameters of the injection are stored in a sealed envelope. Oblivious to this, the rest of the team analyses all events detected as candidates that cross the threshold criteria like signal-to-noise ratio, relative time delay between the detectors, etc. It is only after confirmation or otherwise as an event, that the information whether there was indeed a blind injection would be revealed. After the event GW 150914 went through the preliminary analysis as a candidate, it was confirmed that there was neither a blind injection nor inadvertent or intentional

access to the detector system that could generate the observed signal.

### The binary black hole

The confirmation that the source was a binary black hole (BBH) system with relatively heavy (36 and 29 solar masses) black holes, merged and gone from view for ever now, was indeed a surprise. The detailed arguments for establishing this may be found in references 1 and 11. This discovery indicates a good detectable population of such systems and is sure to excite astrophysicists to reconsider models of black hole formation, especially in binary systems. The observed signal can be reproduced well when such a system is at a distance of about 400 Mpc (signal amplitude is just inversely proportional to the distance, as for electromagnetic waves). It is perhaps important to stress that this is the first observation and confirmation of existence of a stellar mass binary black hole system, adding much value to the already spectacular detection of the waves in a terrestrial GW antenna.

The largely sinusoidal chirp of the waveform indicates that the orbit eccentricity, if any, is small. In highly elliptical orbits, dominant emission is at closest approach with high velocity and results in more spiky nature of the waveform with higher harmonics. Lack of slower periodic modulations of the waveform (except the chirp) puts strong constraints on the magnitude and orientation of the spins, relative to the orbital angular momentum. If the individual black holes have large spins that are not perpendicular to the orbital plane, relativistic precessions will result in modulating the orientation of the orbital plane and will be reflected as modulations of waveform. However, an accurate estimate of spin related parameters requires fairly involved modelling and computations.

### General relativity and the aLIGO event

The general relativistic prediction for the nature and flux of gravitational waves was verified with great accuracy<sup>12</sup> by observing the orbital decay of the Hulse–Taylor binary pulsar system 1513 + 16. The BH–BH gravitational waves observed by aLIGO confirm the GR prediction with

about 10% accuracy in a scenario involving black holes that approach each other close enough for their horizons to overlap, with orbital speed more than 50% of the velocity of light in a region with high gravitational potentials. All the parameters estimated from the waveform are consistent with general relativity. The large orbital angular momentum of the binary system is very well determined from the relativistic Keplerian parameters and from the several cycles of the slow chirp waveform. During the evolution, significant fraction of angular momentum, along with the orbital energy, is radiated in the circularly polarized gravitational waves. The remaining angular momentum and whatever spins the individual black holes go to make the spin of the final Kerr black hole. An estimate shows that the final black hole is spinning at about 70% of the maximum spin that is physically possible<sup>1</sup>. There is very good agreement with what one expects from the general theory of relativity, even though the computationally well determined final spin is not accurately measured directly from the ‘ring-down’ part of the waveform after merger.

The formation of black hole from merger or collapse of matter is not spherically symmetric in general and one expects shaking away of these oscillating quadrupolar modes on the way to the formation of a final spherical black hole, all visible in principle in the emitted gravitational waves after merger. The quasi-periodic oscillations of the forming black hole were investigated in the past, with pioneering work on scattering of gravitational waves from already formed black holes by Vishveswara<sup>13</sup>, and on black hole oscillations by Press<sup>14</sup>, and Chandrasekhar and Detweiler<sup>15</sup>. The large signal from the LIGO event enabled researchers to sieve out some indication of the ring-down towards the final black hole even though the SNR in this region is not high enough for precision tests. In any case, the entire waveform is consistent with GR at about 10% level and this is reflected in similar accuracies of parameters estimated from the waveform using general relativistic calculations.

The parameters of the binary system and the final black hole are well estimated from the detailed analysis of the GW waveform or the relative motion of the mirrors of the interferometers. Considering that the astrophysical ‘event’ happened 1300 million light years away,

this is a great achievement for the first catch in the detectors. The masses are 36 and 29 solar masses (10% accurate), and energy worth 3 solar masses was radiated in waves. At peak emission the power output was  $3.6 \times 10^{49}$  Watts, which is equivalent to 200 suns worth mass converted to energy in a single second! (This level of luminosity lasts only for a few ms, and hence the energy converted at peak is equivalent to less than two solar masses.) That makes the GW150914 event a thousand times more powerful than the brightest gamma-ray bursts. The total energy radiated in gravitational waves (estimated from the observed waveform and estimated distance) is about 5% of the total mass energy of BBH system. It is perhaps important to point out that no matter is directly converted to energy and the entire radiation comes from the gravitational potential energy. After all, these are black holes and no matter can be released from them.

One difference between the optical observation of the decay of binary pulsar and this direct detection is that the orbital decay is sensitive to the total energy radiated away, including hypothetical scalar waves or other possibilities in theories different from general relativity. The detectors on the other hand are designed to be most sensitive to the quadrupolar gravitational waves of transverse nature and for a source position where this sensitivity maximizes, the detectors are fairly insensitive to the scalar waves. Therefore, detailed modelling is required to directly address accurate tests of alternate theories and a rigorous analysis requires starting from exact premises of particular theories.

### The LSC and IndiGO-LSC

The LIGO Scientific Collaboration (LSC) is a group of more than 1000 scientists from more than 90 universities and research institutes around the United States and in 14 other countries including India. The LSC detector network includes the two aLIGO interferometers and the GEO600 detector in Hannover, Germany. Several conceptual and technical developments for the LIGO detectors were contributed by the GEO team that includes scientists at the Max Planck Institute for Gravitational Physics (Albert Einstein Institute, AEI), Leibniz Univer-

sität Hannover, along with partners at the University of Glasgow, Cardiff University, the University of Birmingham, other universities in the United Kingdom, and the University of the Balearic Islands in Spain. There are also significant contributions to LIGO from the Australian Consortium for Interferometric Gravitational Astronomy. In addition, the Virgo collaboration that operates the 3-km Virgo detector in Cascina, Pisa, Italy joins hands to form the larger LIGO–Virgo collaboration and publishes most results jointly.

The current participation of Indian scientists in LSC is under the IndiGO (Indian Initiative for Gravitational Wave Observations) umbrella. The gradually expanding course of gravitational wave research in India, with significant contributions to waveform calculations, data analysis and an impeccable record of collaborative research and training, is imprinted in the aLIGO discoveries and is well-acknowledged in the discovery paper. The IndiGO consortium that originated in 2009 to bind together and take forward GW research in India has now more than 120 members and about half of them are engaged in experimental activities in various fields relevant for GW detectors. About 60 of these are already active members of the LIGO Scientific Collaboration (LSC) and 37 are co-authors in the discovery paper. This includes senior researchers and engineers as well as post-doctoral fellows and graduate students. IndiGO's strong presence in the LSC is due to the commendable and continued history of involvement in theoretical work and data analysis as well as due to the intense work that has been put in to propose and realize the LIGO-India GW detector project, taken up in 2011. This was at a point of realization that there was a significantly large community of GW scientists involved in advanced calculations and data analysis techniques as well as several scientists and engineers keen to direct their expertise in optics and precision metrology towards GW research in India, who could come together and seek support for the construction and operation of an advanced detector in India. The viability of this dream was well-founded on the history of nearly three decades of gravitational physics research in India.

Bala Iyer, who was at RRI, Bengaluru, was a key contributor to precision calcu-

lations<sup>16</sup> of exact shape of the gravitational waves emitted by orbiting astrophysical sources in the 90s. These calculations are so complicated, both conceptually and computationally, that only a few people, spread over two or three groups in the world had ventured into such studies. Recent developments in numerical relativity have of course enormously improved the present scenario. Some pioneering contributions for searching whether there is a genuine signal in the usually noisy data by scanning the data with such characteristic waveforms, looking for the best match, were made by Dhurandhar at IUCAA, Pune and his group<sup>17</sup>. The large theoretical community of GW researchers in India today was seeded by these early researches.

Over the last decade the Indian gravitational-wave community has spread to a number of educational and research institutions in India. These include CMI Chennai, ICTS-TIFR Bangalore, IISER-Kolkata, IISER-Thiruvananthapuram, IIT Gandhinagar, IPR Gandhinagar, IUCAA Pune, RRCAT Indore and TIFR Mumbai. IUCAA and ICTS-TIFR host LIGO Tier-3 grid computing centers. Major contributions of Indian scientists include the development of techniques to coherently combine and analyse the data from multiple observatories, development of hierarchical search methods that allow us to progressively dig into the data, techniques for making sky maps of stochastic gravitational waves, formulating methods to accurately test Einstein's theory of general relativity using gravitational-wave observations, modelling gravitational-wave signals by combining post-Newtonian calculations with large-scale supercomputer simulations, and devising strategies to extract astrophysical information from a joint electromagnetic and gravitational-wave observation of a source.

The gravitational experiments group of TIFR, led by Cowsik, pioneered precision experiments to test gravity theories in India in the mid-eighties with special purpose laboratories and novel instruments<sup>18</sup>. TIFR funded a prototype interferometer GW detector, proposed by IndiGO-LSC members Unnikrishnan and Rajalakshmi in 2011. This new laboratory and prototype instrument are expected to play a significant role in training and research that support the LIGO-India project. The group has contributed

actively to the LIGO-India project proposal.

A number of IndIGO scientists have contributed to the analyses in the double discovery paper to decipher the information about the binary black hole merger event encoded in the detected gravitational waves<sup>19</sup>. The analysis on several fronts confirms that even a most exotic event like the inspiral and merger of two black holes proceeded along the theoretical expectations in general relativity.

### GW detector network

The only aspect that could have made this detection even more spectacular, without doubt, is the identification of the location of the source in the sky. With just 2 detectors we have to be satisfied with a band of possible locations, determined by just one relative delay between the detectors, convolved with the sensitivity and beam pattern of the two detectors. Though such sky bands with location uncertainty of 600 square degrees were circulated to several collaborations that have prior MoU with LSC for electromagnetic follow up, no optical counterpart was detected, understandably. (There is report of a detected transient in X-rays above 50 keV by the Fermi Gamma-ray Burst Monitor, albeit with a delay of 0.4 s and then lasting for a second<sup>20</sup>.) Whether a genuine connection or not, there is much in expectation for the X-ray instruments on board the Indian multi-wavelength astronomy satellite Astrosat. The next big discovery from LIGO is expected to be the identification of an electromagnetic signal (such as a gamma-ray burst) associated with a gravitational wave event. However, this requires a global network of detectors with intercontinental separations to localize gravitational wave events accurately on the sky. Clearly, we need more detectors that can match the exquisite and essential sensitivity of the two aLIGO detectors, with good overlap with their frequency range. The international science community is unanimous that the key to the future of gravitational wave astronomy will be a gravitational wave detector network spread over the globe with capabilities for localizing the source in the sky, which can then be identified with electromagnetic wave telescopes.

The advanced Virgo detector is expected to become operational in 2017,

with good overlap of sensitivity with the aLIGO detectors. This will enable the first phase of gravitational wave astronomy with source location identification with a precision of 10 square degree or so in many parts of the sky. Improvement towards square degree precision requires more aLIGO like detectors and this highlights the importance of early deployment of detectors like LIGO-India. The localization accuracy improves with the separation between detectors. While the baseline is 10 ms, in light travel time, between the US detectors, the maximum baseline possible on the global earth is about 40 ms, and a detector in India will have nearly such a baseline separation from the US detectors. The cryogenic KAGRA underground GW detector<sup>21</sup> below Kamioka mountains, Japan is expected to become online by 2018. The present scenario envisages at least 5 aLIGO level detectors, with slightly different sensitivities and frequency ranges, operational by 2022 and the next decade of astronomy will be aided and indeed might be dominated by GW based astronomy.

### The new wave in astronomy and LIGO-India

Advanced LIGO detectors have been operating in their first phase of observational run (O1) during 14 September 2015 to 14 January 2016. The continuing analysis of the full data will give a good indication of expected rates of such events and it seems clear that the aLIGO detectors at their full sensitivity will regularly detect gravitational waves from sources up to a Gpc, perhaps more than an event per month. This is exciting for gravitational wave astronomy because a good fraction of these sources, involving binary black holes, may not be visible anywhere in the electromagnetic spectrum.

Even this first event in aLIGO detectors proves amply that gravitational wave astronomy has started and has asserted itself as an independent and essential window to the high energy stellar evolution events in the universe. LIGO has opened up a fundamentally new observational window to the Universe. This has the potential to revolutionize our understanding of astrophysics, cosmology and fundamental physics. If the event rate determinations based on the signals so far

turn out to be stable in the long run, one can expect a reliable detection of several events every year in the aLIGO detectors operating at their full sensitivity<sup>22</sup>. Though the BH–BH type of events may not be visible to electromagnetic telescopes, events involving binary neutron stars are expected to show associated signals in telescopes spanning the entire electromagnetic spectrum. However, source localization is the key to successful follow-ups and integrated multi-wavelength multi-wave astronomy and this is indeed the future focus.

One of the key installations to realize a network with aLIGO capabilities is the ambitious LIGO-India project<sup>23,24</sup>. Proposed in 2011, this project aims to build an Advanced LIGO detector on Indian soil in collaboration with LIGO-USA and its international partners Germany, UK and Australia. Expected to start joint observation with the US detectors by 2023, LIGO-India will be the first international frontier science experiment on Indian soil, and will involve cutting-edge technology in lasers, optics, ultra-high vacuum and control system engineering. The project will bring together the best of fundamental science and high-end technology available in the country at national research laboratories, IITs, IISERs, universities and industry. LIGO-India will involve huge Indian industry involvement. The Department of Atomic Energy (DAE) and the Department of Science and Technology (DST) have joined hands to support this megascience project. Almost the entire cost of the project will be expended in industries and laboratories within the country. It would serve as an internationally visible flagship for Industry–Academia partnership. Lead institutes for the proposal are IUCAA, IPR and RRCAT, in collaboration with the IndIGO consortium. IUCAA is responsible for site selection (IISER-Kolkata contributing in seismic characterization), data analysis and computing facilities, science and human resource development; IPR for civil infrastructure and facilities, vacuum system and mechanical engineering; RRCAT for the optics, detector integration, installation and commissioning. Teams at the three lead institutions have been intensely involved in taking forward all initial tasks related to the project. The proposal was cleared by the Union cabinet soon after the announcement of the discovery in February 2016

with an ‘in-principle approval’ for the LIGO-India project and this will give the much needed big boost, public support and operational funds to the LIGO-India detector project.

1. Abbott, B. P. *et al.*, *Phys. Rev. Lett.*, 2016, **116**, 061102.
2. Saulson, P., *Fundamentals of Interferometric Gravitational Wave Detectors*, World Scientific, Singapore, 1994.
3. Aasi, J. *et al.*, *Class. Quantum Grav.*, 2015, **32**, 074001.
4. Abbott, B. P. *et al.*, *The Advanced LIGO Detectors in the Era of First Discoveries*, 2016; arXiv:1602.03838
5. Weiss, R., Electromagnetically coupled broadband gravitational antenna, Tech. Rep., Quarterly report of the Research Laboratory for Electronics, MIT, 1972.
6. Moss, G. E., Miller, L. R. and Forward, R. L., *Appl. Opt.*, 1971, **10**, 2495–2498.
7. Forward, R. L., *Phys. Rev.*, 1978, **17**, 379–390.
8. Drever, R. W. P., Raab, F. J., Thorne, K. S., Vogt, R. and Weiss, R., A Laser Interferometer Gravitational-wave Observatory (LIGO), Technical Report, 1989.
9. Brilliet, A. *et al.*, The Virgo Project, Technical Report, VIR-0517A-15, Virgo Collaboration, 1989.
10. LIGO Scientific Collaboration, LIGO-P1500238, 2016.
11. Abbott, B. P. *et al.*, LIGO scientific collaboration and Virgo collaboration, Properties of the binary black hole merger GW150914, 2016; arXiv:1602.03840.
12. Taylor, J. H. and Weisberg, J. M., *Astrophys. J.*, 1982, **253**, 908–920.
13. Vishveshwara, C. V., *Nature*, 1970, **227**, 936–938.
14. Press, W. H., *Astrophys. J.*, 1971, **170**, L105–L108.
15. Chandrasekhar, S. and Detweiler, S. L., *Proc. R. Soc. London*, 1975, **A344**, 441–452.
16. Blanchet, L. *et al.*, *Phys. Rev. Lett.*, 1995, **74**, 3515–3518.
17. Sathyaprakash, B. and Dhurandhar, S., *Phys. Rev.*, 1991, **D44**, 3819–3834.
18. Unnikrishnan, C. S., *Class. Quantum Grav.*, 1994, **11**, A195–A206.
19. Details of press releases are available at gw.iucaa.in
20. Connaughton, V. *et al.*, (Fermi Collaboration), Fermi GBM Observations of LIGO Gravitational Wave event GW150914, 2016; arXiv:1602.03920.
21. Aso, Y. *et al.*, *Phys. Rev.*, 2013, **D88**, 043007.
22. Abbott, B. P. *et al.*, LIGO scientific collaboration and Virgo collaboration, The rate of binary black hole mergers inferred from advanced LIGO observations surrounding GW150914, 2016; arXiv:1602.03842.
23. Iyer, B. *et al.*, LIGO-India, Technical Report LIGOM1100296, 2011; <https://dcc.ligo.org/LIGO-M1100296/public/main>
24. Unnikrishnan, C. S., *Int. J. Mod. Phys.*, 2013, **D22**, 1341010.

ACKNOWLEDGEMENTS. This article was compiled with published information from the LSC in the detection paper and some of the companion papers, the LSC and IndIGO press releases. Updates and details can be consulted at the IndIGO website [www.gw-indigo.org](http://www.gw-indigo.org) and the LSC website [www.ligo.org](http://www.ligo.org).

---

*C. S. Unnikrishnan is in the Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India. e-mail: unni@tifr.res.in*