Presence of glauconite and kaolinite has additionally been observed at this locality. The evidence is suggestive of the source of sediments to lie in the vicinity in the near shore areas.

Acknowledgements. The authors wish to accord their gratitude to the competent authority of ONGC for granting permission to publish this paper. We are grateful to Shri.P.S.N.Kutty, GGM-Basin Manager, Cauvery Basin, ONGC, Chennai for permission to carry out this work. We wish to acknowledge Dr.D.S.N.Raju, Retd. DGM (Geol), ONGC for his encouragement for this work. Thanks are due to Shri B. Bhaskaran, Suptdg.Geologist for his assistance in labotratory analysis. S/Shri K.Parthiban, R.Parthiban and Selva Raj are acknowledged for their help in collection of samples. The views expressed are those of the authors and not necessarily of the organisation to which they belong.

References

Heinemann Educational Books, London, 244p.

- BARKER, R.W. (1960) Taxonomic Notes on the species figured by H.B.Brady in his report on the Foraminifera dredged by H.M.S.Challenger during the years 1873-1876e, SEPM Spl. Publ. No.9, USA, 115p.
- LOEBLICH, A.R. Jr. and TAPPAN, H. (1988) Foraminiferal genera and their classification. Van Nostrand Reinhold, New York, 869p.

MURRAY, J.W. (1971) An Atlas of British Recent Foraminiferids.

- MURRAY, J.W. (1973) Distribution and ecology of living benthic foraminiferids, Crane, Russak and Company Inc., 274p.
- MURRAY, J.W. (1991) Ecology and paleoecology of benthic foraminifera, Longman, Essex, 397p
- WHITTAKER, J.E and HODGKINSON, R.L. (1979) Foraminifera of the Togopi Formation, Eastern Sabah, Malaysia, Bull. British Museum, Geol. Series, v.31, no.1, pp 1-120.

(Received: 24 August 2005; Revised form accepted: 7 July 2006)



JOUR.GEOL.SOC.INDIA, VOL.69, JAN. 2007

74