



ABS049

ASSESSMENT OF 1P/19Q DELETION BY FLOURESCENCE INSITU HYBRIDIZATION (FISH) IN GLIOMA PATIENTS FROM A SINGLE INSTITUTE EXPERIENCE

**Karthika Ashok
Kumar¹,**

Akhila Sivadas²,
Ananya Dutt AS³,
Chanshi Chandran⁴,
M.V Thampi⁵,
Pavithran K⁶

¹Pharmd Final Year
Amrita School Of
Pharmacy, Kochi

²Clinical Pharmacist
Department of Oncology,
Amrita Institute of
Medical Sciences, Kochi

³Pharm D Intern Amrita
School of Pharmacy,
Kochi

⁴Pharm D Final Year
Amrita School of
Pharmacy

⁵Department of
Pathology, Amrita
Institute of Medical
Sciences, Kochi

⁶Head of the Department
Medical Oncology &
Hematology, Amrita
Institute of Medical
Sciences. Kochi

The discovery of 1p and 19q chromosomal arms deletion in glial tumors influences both more objective diagnosis and more accurate prediction of chemotherapy response.

As a result an attempt has been made to detect deletion using fluorescence in-situ hybridization (FISH) and analyzed its prognostic value in a cohort of glial tumor patients from Amrita Institute of Medical Sciences and Research Center Kochi .

FISH was performed on 66 FFPE tissue sections by using Vyis LSI 1p36/LSI 1q25 and LSI 19p13/LSI 19q13 dual coloured FISH probe sets. Signals were scored from at least 150-250 non-overlapping, intact nuclei. 163 cases were analyzed. Both 1p and 19q deletions was observed only in 28/163 (17.17%) , - 1p/+ 19q deletion 80/163 (49.07%) and +1p/-19q deletion 55/163(33.74%) .

In this work presented the FISH was successfully applied to identify deletion 1p/19q. Its incidence depends on the type of diagnosed gliomas. In contrast to reported data, the present study reveals 49.07% deletion - 1p/+ 19q

Deletions also have prognostic significance in the test group what constitutes the basis for inclusion of determining deletion 1p/19q into diagnostic and treatment algorithm.

Keywords: FISH, deletion 1p/19q, glioma