



An updated Checklist of Superfamily Drepanoidea (Lepidoptera) from India

Rahul Joshi^{1*}, Navneet Singh², Gyula M. László³ and Jalil Ahmad²

¹Zoological Survey of India, GPRC, Sector-8, Bahadurpur Housing Colony, Patna - 800026, Bihar, India;
Email: joshiarctiidae@gmail.com

²Lepidoptera section, Zoological Survey of India, New Alipore, Kolkata - 700053, West Bengal, India;
Email: nsgill007@gmail.com; jalilahmad23046@gmail.com

³The African Natural History Research Trust (ANHRT), Street Court Leominster-Kingsland, HR6 9QA, United Kingdom; Email: gyula.m.laszlo@gmail.com

Abstract

An updated checklist of 164 valid species (including subspecies) under 55 genera of superfamily Drepanoidea, family Drepanidae representing four subfamilies: Cyclidiinae, Drepaninae, Oretinae and Thyatirinae has been compiled. The detailed information about distribution within India as well as in other countries, first reference, synonymy has been provided for each species. Clarifications regarding distributional limits within India are also given.

Keywords: Drepanidae, Cyclidiinae, Drepaninae, Oretinae and Thyatirinae

Introduction

Superfamily Drepanoidea is a member of clade Macroheterocera (Glossata: Lepidoptera). The classification of the superfamily Drepanoidea and the position of its members remained ambiguous during the last century. Hampson (1892) studied Drepanoidea as Drepanulidae; Imms (1934) placed the members of Drepanoidea in the Geometroidea and later studies (McDunnough, 1938; Nakamura, 1981; Minet, 1983, 1991; Scoble, 1992; Holloway, 1998; Minet and Scoble, 1999; Holloway et al., 2001; Kristensen *et al.*, 2007; Regier *et al.*, 2009 and Mutanen *et al.*, 2010; Wu *et al.*, 2010) redefined the superfamily towards its stability. At present, Drepanoidea includes three families: Cimeliidae, Doidae and Drepanidae with an estimate of 672 described species from the Globe (Nieukerken *et al.*, 2011). As far as Indian fauna is concerned, Drepanoidea is represented by the only family, Drepanidae with no consolidated information on its species composition. First important compilation of Indian Drepanidae was attempted by Hampson (1892)

(as Drepanulidae) with inclusion of 66 species from then limits of the British India. Family Drepanidae is defined by their characteristic tympanal organs derived from tergo-sternal sclerites connecting sternum A₂ with tergum A₁, and can be seen as a pair of small oval structures, each composed of two interconnecting chambers; frenulum, when present, is often apically clubbed; forewing venation often shows reduction of branches of the radial system, usually with a narrow areole within it; genitalia of both sexes show variety of structure, tegumen usually with socii; aedeagus vesica rarely has cornuti; female genitalia often has a signum or signa (Holloway, 1998). Wu *et al.* (2010) restored subfamily Oretinae and divided Drepanidae into four subfamilies: Cyclidiinae, Drepaninae, Oretinae and Thyatirinae. Present paper follows the same classification. In addition to above mentioned works, other publications on the species composition of Indian Drepanidae or with representation of Indian species in their studies are as follows: Hampson (1893, 1895, 1896, 1912, 1914), Holloway (2011), Warren (1896, 1897, 1901, 1902, 1914, 1922, 1923), Swinhoe (1899), Watson (1957, 1957a, 1959, 1961, 1965, 1967, 1968),

* Author for correspondence

Walker (1860, 1862, 1862a, 1866), Wilkinson (1967, 1968, 1970, 1972), Yazaki (1992, 1993), László *et al.* (2001, 2007), Werny (1966) and Yoshimoto (1982, 1987, 1990, 1992, 1993), Chandra & Sambath (2013), Jiang *et al.* (2015, 2016), Zhuang *et al.*, (2017) and Sanyal *et al.*, 2018).

The present paper provides the checklist of 164 valid species and subspecies under 55 genera of family Drepanidae from India covering all the four subfamilies: Cyclidiinae, Drepaninae, Oretinae and Thyatirinae. The family is well distributed in Indian region representing about 1/4th of the known species of the world. The information presented in this checklist is compiled from published literature as mentioned above and as mentioned under distribution of each species. For each species, updated nomenclature, first reference, synonymy, distribution within India as well as other part of the Globe is also provided. Synonymised species added under species has been updated from the references mentioned under distribution of that species. Information about the sister subspecies of the taxa dealt here and some errors in earlier published literature regarding distribution within India as well as other related information has been updated and mentioned under remarks.

Systematic Account

Superfamily DREPANOIDEA

Family DREPANIDAE

Subfamily CYCLIDIINAE

Genus *Cyclidia* Guenée, 1857

Guenée, 1857; *Hist. nat. Ins., Spec. gén. Lépid.*, 9: 62.

Type-species: *Euchera substigmata* Hübner, 1831.

1. *Cyclidia dictyaria* (Swinhoe, 1899)

1899. *Euchera dictyaria* Swinhoe, *Ann. Mag. Nat. Hist.*, 7(3): 111.

1922. *Cyclidia dictyaria*: Warren, in Seitz, *Gross-Schmett. Erde*, 10: 445.

Distribution: India: N.W. India, Kangra (Swinhoe, 1899); Karnataka (Warren, 1922). *Elsewhere*: not known.

2. *Cyclidia substigmata superstigmata* Prout, 1918

1918. *Cyclidia superstigmata* Prout, *Ann. Mag. Nat. Hist.*, 9(2): 416.

2016. *Cyclidia substigmata superstigmata*: Jiang *et al.*, *Zookeys*, 553: 125.

Distribution: India: Dharamshala, Assam, Sikkim (Hampson, 1892); Darjeeling (Jiang *et al.*, 2016). *Elsewhere*: Nepal, Burma (Hampson, 1892); Nepal (Jiang *et al.*, 2016).

Remarks: Jiang *et al.* (2016) considered five valid subspecies under *Cyclidia substigmata* (Hübner, 1831) and treated *Cyclidia substigmata brunnea* Chu & Wang, 1987 and *Cyclidia tetraspota* Chu & Wang, 1987 as its synonym. The nominotypical subspecies from China and Vietnam; *C. substigmata intermedia* Prout, 1918 (Tibet); *C. substigmata nigralba* Warren, 1914 (Japan and Korea); *C. substigmata modesta* Bryk, 1943 (Mynamar) are not known from India.

3. *Cyclidia rectificata rectificata* (Walker, 1862)

1862. *Nelcynda rectificata* Walker, *List Lepid. Het. Br. Mus.*, 24: 1142.

1862a. *Cyclidia muricolaria* Walker, *ibid.*, 25: 1483.

1866. *Cyclidia patulata* Walker, *ibid.*, 35: 1537.

2016. *Cyclidia rectificata*: Jiang *et al.*, *Zookeys*, 553: 130.

Distribution: India: Sikkim, Nagaland (Hampson, 1892); Shillong (Hampson, 1895); Arunachal Pradesh (Chandra & Sambath, 2013); Darjeeling, Sikkim (Jiang *et al.*, 2016). *Elsewhere*: China, Tibet (Jiang *et al.*, 2016).

Remarks: The second subspecies, *C. rectificata malaisei* Bryk, 1943 is reported from Myanmar.

Subfamily DREPANINAE

Genus *Agnidra* Moore, 1867

Moore, 1867; *Proc. Zoo. Soc. Lond.*, 1867: 618.

Type-species: *Fascellina specularia* Walker, 1866.

4. *Agnidra specularia* (Walker, 1866)

1866. *Fascellina specularia* Walker, *List Lepid. Het. Br. Mus.*, 35: 1553.

1943. *Albara ochracina* Bryk, *Ark. Zool.*, 34(A13): 17.

1968. *Agnidra specularia*: Watson, *Bull. Br. Mus. nat. Hist. Suppl.*, 12: 36.

Distribution: India: N.E. India (Darjeeling, Khasia Hills, Cherrapunji), Sikkim (Watson, 1968). *Elsewhere*: Bhutan, Burma, Ceylon and Vietnam (Watson, 1968), Nepal (Wilkinson, 1972; Yazaki, 1992).

5. *Agnidra corticata corticata* (Warren, 1922)

1922. *Drepana corticata* Warren, in Seitz, *Gross-Schmett. Erde*, 10: 464.

1968. *Agnidra corticata corticata*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.) Suppl.*, 12: 39.

Distribution: India: NE India (Darjeeling, Khasis) (Watson, 1968). *Elsewhere*: not known.

Remarks: Its another subspecies, *Agnidra corticata francki* Watson, 1968 reported from China.

6. *Agnidra vinacea* (Moore, 1879)

1879. *Drepana vinacea* Moore, *Descr. Indian lepid. Atkinson*, (1): 85.
 1943. *Albara birmanica* Bryk, *Ark. Zool.*, **34A** (13): 18.
 1968. *Agnidra vinacea*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.) Suppl.*, **12**: 40.

Distribution: India: Sikkim, Khasis, Naga Hills, N.E. India (Watson, 1968); Uttarakhand (Dey & Sondhi, 2019). *Elsewhere*: N.E. Burma (Watson, 1968); Nepal (Wilkinson, 1972).

7. *Agnidra discipilaria* Moore, 1868

1868. *Agnidra discipilaria* Moore, *Proc. Zool. Soc. Lond.*, **1867**: 619.
 1886. *Agnidra usta* Butler, *Illustr. Lep. Het. Br. Mus.*, **6**: 17.
 1922. *Albara magnidiscata* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 468
 1943. *Albara discipilaria macularis* Bryk, *Ark. Zool.*, **34A** (13): 18
 1968. *Agnidra discipilaria*: Watson, *Bull. Br. Mus. Nat. Hist. Suppl.*, **12**: 44.

Distribution: India: N.E. India, Darjeeling (Gopaldhara), Khasis, Sikkim (Watson, 1968); Uttarakhand (Dey & Sondhi, 2019). *Elsewhere*: Burma and Thailand (Watson, 1968); Nepal (Wilkinson, 1972)

Genus *Albara* Walker, 1866

Walker, 1866; *List Lep. Het. Br. Mus.*, **35**: 1566.

Type-species: *Albara reversaria* Walker, 1866.

8. *Albara reversaria* Walker, 1866

1866. *Albara reversaria* Walker, *List Lep. Het. Br. Mus.*, **35**: 1567.
 1897. *Albara opalescens* Warren, *Novit. Zool.*, **4**: 12.
 1914. *Albara griseotincta* Wileman, *Entomologist*, **47**: 268.
 1921. *Albara horishana* Matsumura, *Thousand Ins. Japan Suppl.*, **4**: 948.
 1998. *Albara reversaria*: Holloway, *Moths of Borneo*, **8**: 56.

Distribution: India: N.E. Himalaya, Khasis (Holloway, 1998). *Elsewhere*: Sumatra, Borneo, Peninsular Malaysia, Taiwan (Holloway, 1998).

Genus *Auzata* Walker [1863]

Walker [1863]; *List Lep. Het. Br. Mus.*, **26**: 1620.

Type-species: *Auzata semipavonaria* Walker, [1863].

9. *Auzata ocellata* (Warren, 1896)

1896. *Gonocilix ocellata* Warren, *Novit. Zool.*, **3**: 337.
 1897. *Auzata ocellata*: Hampson, *J. Bomb. Nat. His. Soc.*, **11**: 287.

Distribution: India: Assam, Khasis (Watson, 1959). *Elsewhere*: Burma, China (Watson, 1959).

10. *Auzata semipavonaria* Walker, [1863]

1863. *Auzata semipavonaria* Walker, [1863]; *List Lep. Het. Br. Mus.*, **26**: 1620.

Distribution: India: Mussoorie, Shimla, Kullu, Sikkim, Pakistan (Rawalpindi) (Hampson, 1892; Watson, 1959). *Elsewhere*: Pakistan (Rawalpindi) (Watson, 1959).

11. *Auzata simplicata* Warren, 1897

1897. *Auzata simplicata* Warren, *Novit. zool.*, **4**: 13.

Distribution: India: North India (Mussoorie, Darjeeling, Nainital, Khasis) (Watson, 1959). *Elsewhere*: China (Watson, 1959)

Genus *Auzatellodes* Strand, 1917

Strand, 1917; *Arch. Naturgesch.*, **82** A(3): 149.

Type-species: *Auzatellodes desquamata* Strand, 1917.

12. *Auzatellodes hyalinata* (Moore, 1868)

1868. *Comibaena hyalinata* Moore, *Proc. Zool. Soc. Lond.*, **1867**: 638.
 1943. *Hyalospectra postfasciata* Bryk, *Ark. Zool.*, **34**(A 13): 7.
 1988. *Auzatellodes hyalinata*: Inoue, 1988; *Tinea*, **12**: 125-134.

Distribution: India: North-East Himalaya, NE Bengal, Sikkim (Inoue, 1988). *Elsewhere*: Nepal, North Burma, China, Sri Lanka, Thailand (Wilkinson, 1972).

Genus *Callidrepana* Felder, 1861

Felder, 1861; *S.B. Akad. Wiss. Wien*, **43**(1): 30

Type-species: *Callidrepana saucia* Felder, 1861.

13. *Callidrepana argenteola* (Moore, 1858)

1858. *Drepana argenteola* Moore, *Cat. Lepid. Insects Mus. Hon. E. India Co.*, **2**: 369.
 1893. *Drepana bracteata* Hampson, *Illustr. Lep. Het. Br. Mus.*, **9**: 68.
 1922. *Callidrepana bracteata celebensis* Warren, In Seitz, *Gross-Schmett. Erde*, **10**: 471.
 1932. *Callidrepana dialitha* West, *Novit. Zool.*, **37**: 228.
 1998. *Callidrepana argenteola*: Holloway, 1998; *Moths of Borneo*, **8**: 63.

Distribution: India: Sikkim, Burma, Sri Lanka, Java (Hampson, 1892), Uttarakhand (Smetacek, 2002). *Elsewhere*: Oriental tropics to Taiwan, Philippines, Sulawesi and Timor (Holloway, 1998).

14. *Callidrepana patrana patrana* (Moore, 1866)

1866. *Drepana patrana* Moore, *Proc. Zool. Soc. Lond.* **1865**(3): 816.
 1943. *Callidrepana patrana subbasalis* Bryk, *Ark. Zool.*, **34**(A 13): 21.
 1955. *Callidrepana patrana formosana* Inoue, *Butterfl. Moths*, **6**: 13.

1968. *Callidrepana patrana patrana*: Watson, 1968, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 116.

Distribution: India: NE India (Darjeeling) (Watson, 1968). *Elsewhere*: North Burma, Taiwan, China (Watson, 1968); Nepal (Wilkinson, 1972).

Remarks: Its sister subspecies, *Callidrepana patrana palleolus* (Motschulsky, 1866) reported from Japan.

15. *Callidrepana gemina gemina* Watson, 1968

1968. *Callidrepana gemina gemina* Watson, *Bull. Br. Mus. Nat. Hist. (Ent.) Suppl.*, **12**: 119.

Distribution: India: NE India, Darjeeling (Watson, 1968). *Elsewhere*: not known.

Remarks: *Callidrepana gemina curta* Watson, 1968 is known from China.

16. *Callidrepana gelidata* (Walker, 1862)

1862. *Damna gelidata* Walker, *List Lep. Het. Br. Mus.*, **26**: 1570.

1889. *Platypteryx argentilinea* Snellen, *Tijdschr. Ent.*, **32**: 8.

1998. *Callidrepana gelidata*: Holloway, *Moths of Borneo*, **8**: 62.

Distribution: India: Location not specify (Holloway, 1998). *Elsewhere*: Borneo, Peninsular Malaysia, Singapore, Sumatra (ZSM), Java, Burma (Holloway, 1998).

17. *Callidrepana micacea* (Walker, 1862)

1862. *Drepana micacea* Walker, *J. Linn. Soc. Lond. (Zool.)*, **6**: 176.

1879. *Drepana biocularis* Moore, *Descr. Indian lepid. Atkinson*, **1**: 84.

1897. *Ausaris fulvilauta* Warren, *Novit. Zool.*, **4**: 12.

1905. *Callidrepana pilana* Swinhoe, *Ann. Mag. nat. Hist.* **15**(7): 150.

1998. *Callidrepana micacea*: Holloway, 1998; *Moths of Borneo*, **8**: 64.

Distribution: India: N.E. Himalaya (Holloway, 1998). *Elsewhere*: Andamans, Peninsular Malaysia, Sumatra (ZSM), Borneo, Philippines, Sulawesi (Holloway, 1998).

18. *Callidrepana obliquistriga* Warren, 1922

1922. *Callidrepana obliquistriga* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 471.

Distribution: India: Assam (Warren, 1922). *Elsewhere*: not known.

Genus *Cancucha* Walker, 1866

Walker, 1866; *List Lep. Het. Br. Mus.*, **35**: 1574.

Type-species: *Canucha curvaria* Walker, 1866.

19. *Canucha specularis* (Moore, 1879)

1879. *Drepana specularis* Moore, *Proc. Zool. Soc. Lond.*, **1879**: 4077.

1900. *Platypteryx obruncata* Warren, *Novit. Zool.*, **7**: 117.

1968. *Canucha specularis*: Watson, 1968; *Bull. Br. Mus. Nat. Hist. Suppl.*, **12**: 99.

Distribution: India: Sikkim (Watson, 1968). *Elsewhere*: China, Java, Sumatra, Borneo, Sri Lanka, Sulawesi (Watson, 1968), Indian Subregion, S. China, Sundaland, Sulawesi (Holloway, 1998).

20. *Canucha duplexa duplexa* (Moore, 1866)

1866. *Drepana duplexa* Moore, *Proc. Zool. Soc. Lond.*, **1865** (3): 816.

1968. *Canucha duplexa duplexa*: Watson, 1968; *Bull. Br. Mus. nat. Hist. Suppl.*, **12**: 102.

Distribution: India: NE India, Darjeeling (Watson, 1968). *Elsewhere*: not known.

Remarks: Its sister subspecies, *Canucha duplexa birmana* Bryk, 1943 is known from North East Burma.

21. *Canucha miranda* Warren, 1922

1922. *Canucha miranda* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 475.

1931. *Canucha miranda formosicola* Matsumura, 6000 *Illust. Insects Japan.-Empire*, **1931**: 741.

1968. *Canucha miranda*: Watson, *Bull. Br. Mus. Nat. Hist. Suppl.*, **12**: 103.

Distribution: India: Assam (Khasis), N.E. India (Watson, 1968). *Elsewhere*: Formosa (Watson, 1968).

Genus *Cilix* Leach, [1815]

Leach, [1815]; In *Brewster, Edinburgh Ency.*, **9**: 135.

Type-species: *Bombyx compressa* Fabricius, 1777.

22. *Cilix glaucata* (Scopoli, 1763)

1763. *Phalaena glaucata* Scopoli, *Ent. Carniolica*, **1763**: 221.

1767. *Bombyx ruffa* Linnaeus, *Syst. Nat.*, **2**: 1068.

1777. *Bombyx compressa* Fabricius, *Genera Insectorum*, **1777**: 279.

1994. *Cilix glaucata*: de Freina, *Atlanta*, **25**(1/2): 340.

Distribution: India: Shimla, Dalhousie (de Freina, 1994). *Elsewhere*: North America, Europe (Hampson, 1892), Asia monor (de Freina, 1994).

Genus *Deroca* Walker, 1855

Walker, 1855; *List Lep. Het. Br. Mus.*, **4**: 822.

Type-species: *Deroca hyalina* Walker, 1855.

23. *Deroca hidda bifida* Watson, 1957

1957. *Deroca hidda bifida* Watson, *J. Nat. Hist.*, **12**(10): 137.

Distribution: India: Manipur, Naga Hills, Sikkim (Watson, 1957). *Elsewhere:* China (Watson, 1957a).

Remarks: The similar geographical pattern mentioned by Watson (1957) along with the genital differences with nominate subspecies are quite distinct and the species status may be revised in future studies after examination of Holotypes.

24. *Deroca hidda hidda* Swinhoe, 1900

1900. *Deroca hidda* Swinhoe, *Ann. Mag. Nat. Hist.*, 7(6): 306.

1911. *Deroca hyalina subhyalina* Strand, *Gross-Schmetterl.*, 2: 203.

1957. *Deroca hidda hidda*: Watson, *J. Nat. Hist.*, 12(10): 135.

Distribution: India: Jaintia and Khasis Hills (Watson, 1957a). *Elsewhere:* Burma (Watson, 1957a).

25. *Deroca hyalina hyalina* Walker, 1855

1855. *Deroca hyalina* Walker, *List Lep. Het. Br. Mus.*, 4: 823.

1957. *Deroca hyalina hyalina*: Watson, *J. Nat. Hist.*, 12(10): 132-133.

Distribution: India: Dharmshala, Mussorie, Kumaon, Kullu, Kangra, Shimla, NE Himalayas, Manipur, Karo Hills (Watson, 1957a). *Elsewhere:* Burma (Watson, 1957a).

Remarks: *Deroca hyalina latizona* Watson, 1957 is known from China.

26. *Deroca inconclusa concinna* Warren, 1922

1922. *Deroco concinna* Warren, in Seitz, *Gross-Schmett. Erde*, 10: 448.

1957. *Deroca inconclusa concinna*: Watson, *Ann. Mag. Nat. Hist.*, 12(10): 143.

Distribution: India: North East India, Nagaland, Manipur (Watson, 1957a).

27. *Deroca inconclusa inconclusa* (Walker, 1856)

1856. *Cypra inconclusa* Walker, *List Lep. Het. Br. Mus.*, 7: 1727.

1872. *Deroca maculata* Moore, *Proc. Zoo. Soc. Lond.*, 1872: 575.

1957. *Deroca inconclusa inconclusa*: Watson, *Ann. Mag. Nat. Hist.*, 12(10): 140.

Distribution: India: N.W. Himalayas, Mussorie, Dalhousie, Sikkim (Watson, 1957a). *Elsewhere:* Burma, China, Taiwan, Korea, Japan (Hampson, 1892; Watson, 1957a).

Remark: Two more subspecies, *Deroca inconclusa carinata* Watson, 1957 (China) and *Deroca inconclusa phasma* Butler, 1878 (Taiwan, Japan) are not known from India.

Genus *Dipriodonta* Warren, 1897

Warren, 1897; *Novit. Zool.*, 4: 14.

Type-species: *Dipriodonta sericea* Warren, 1897.

28. *Dipriodonta sericea* Warren, 1897

1897. *Dipriodonta sericea* Warren, *Novit. Zool.*, 4: 14.

Distribution: India: Khasi Hills (Wilkinson, 1970). *Elsewhere:* not known.

Genus *Ditrigona* Moore, 1888

Moore, 1888; *Descr. Indian lep. Atkinson* (3): 258.

Type-species: *Urapteryx triangularia* Moore, [1868]

29. *Ditrigona diana* Wilkinson, 1968

1968. *Ditrigona diana* Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 420-421.

Distribution: India: NE India (Darjeeling, Shillong, Naga hills) (Wilkinson, 1968). *Elsewhere:* Bhutan, Nepal (Wilkinson, 1972)

30. *Ditrigona fasciata* (Hampson, 1892)

1892. *Drepana fasciata* Hampson, *Fauna Br. India (Moths)*, 1: 335.

1968. *Ditrigona fasciata*: Wilkinson, *Trans. Soc. Zoo. Lond.*, 31: 459.

Distribution: India: Sikkim (Wilkinson, 1968). *Elsewhere:* not known.

31. *Ditrigona furvicosta* (Hampson, 1912)

1912. *Leucodrepanu furvicosta* Hampson, *J. Bombay nat. Hist. Soc.*, 21 (4): 1271.

1968. *Ditrigona furvicosta*: Wilkinson, 1968; *Trans. Soc. Zoo. Lond.*, 31: 428.

Distribution: India: Sikkim, Darjeeling (Wilkinson, 1968). *Elsewhere:* West China, Tibet, Nepal (Wilkinson, 1972).

32. *Ditrigona inconspicua* (Leech, 1898)

1898. *Teldenia inconspicua* Leech, *Trans. Ent. Soc. Lond.*, 1898: 363.

1968. *Ditrigona inconspicua*: Wilkinson, *Trans. Soc. Zoo. Lond.*, 31: 425.

Distribution: India: Khasi Hills (Warren, 1922). *Elsewhere:* West China (Wilkinson, 1968).

Remarks: Although Warren (1922) mentioned examples of this species from Khasi Hills in Tring museum. Wilkinson (1968) restricted its distribution to China only on the basis of specimen present in British Museum of Natural History.

33. *Ditrigona idaeoides* (Hampson, 1892)

1892. *Leucodrepana idaeoides* Hampson, *Fauna Br. India (Moths)*, 1: 33.

1968. *Ditrigona idaeoides*: Wilkinson, *Trans. Soc. Zoo. Lond.*, 31: 446.

Distribution: India: Sikkim, Darjeeling (Wilkinson, 1968); Uttarakhand (Smetacek, 2002). *Elsewhere:* China (Wilkinson, 1968).

34. *Ditrigona innotata* (Hampson 1892)

1892. *Drepana innotata* Hampson *Fauna Br. India (Moths)*, 1: 335.
1968. *Ditrigona innotata*: Wilkinson, *Trans. Soc. Zool. Lond.*, 31: 433.

Distribution: India: Kullu (Wilkinson, 1968); Uttarakhand (Smetacek, 2002). *Elsewhere:* China (Wilkinson, 1968).

35. *Ditrigona marmorea* Wilkinson, 1968

1968. *Ditrigona marmorea* Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 471-472.

Distribution: India: Assam (Mishmi Hills) (Wilkinson, 1968). *Elsewhere:* China (Wilkinson, 1968).

36. *Ditrigona mytylata* (Guenee, 1857)

1857. *Corycia mytylata* Guenee, *Spec. gen. Ins., Uraniides et Phalenes*, 10(2): 58.
1968. *Ditrigona mytylata*: Wilkinson, *Trans. Soc. Zool. Lond.*, 31: 481.

Distribution: India: North India (Darjeeling, Mussoorie) (Wilkinson, 1968). *Elsewhere:* Burma, Nepal (Wilkinson, 1972).

37. *Ditrigona obliquilinea obliquilinea* (Hampson, 1892)

1892. *Leucodrepana obliquilinea* Hampson, *Fauna Br. India (Moths)*, 1: 333.
1968. *Ditrigona obliquilinea obliquilinea*: Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 445.

Distribution: India: Assam, Sikkim (Wilkinson, 1968). *Elsewhere:* Nepal, Burma (Wilkinson, 1972).

Remarks: *Ditrigona obliquilinea thibetaria* (Poujade, 1895) and its synonym *Corycia pnocaria* Oberthur, 1923 are known from China and Tibet.

38. *Ditrigona pruinosa* (Moore, 1888)

1888. *Thaleridia pruinosa* Moore, *Descr. Indian lepid. Atkinson*, 1888: 261.
1968. *Ditrigona pruinosa*: Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 422.

Distribution: India: Darjeeling (Wilkinson, 1968). *Elsewhere:* not known.

39. *Ditrigona quinaria quinaria* (Moore, 1867)

1867. *Drepanodes quinaria* Moore, *Proc. Zool. Soc. Lond.*, 1867: 618.
1968. *Ditrigona quinaria quinaria*: Wilkinson, 1968; *Trans. Zool. Soc. Lond.*, 31: 441.

Distribution: India: Darjeeling (Wilkinson, 1968). *Elsewhere:* Tibet (Wilkinson, 1968).

Remarks: Apart from nominate and below mentioned subspecies, three other subspecies are also known: *Ditrigona quinaria erminea* Wilkinson, 1968 and *Ditrigona quinaria spodia* Wilkinson, 1968 from China; *Ditrigona quinaria leucophaea* Wilkinson, 1968 from Tibet.

40. *Ditrigona quinaria nivea* (Hampson, 1892)

1892. *Leucodrepana nivea* Hampson, *Fauna Br. India (Moths)*, 1: 333.
1968. *Ditrigona quinaria nivea*: Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 441.

Distribution: India: Khasis, Shillong (Wilkinson, 1968). *Elsewhere:* not known.

41. *Ditrigona regularis* Warren, 1922

1922. *Ditrigona regularis* Warren, in Seitz, *Gross-Schmett. Erde*, 10: 463.
1943. *Ditrigona regularis differentiata* Bryk, *Ark. Zool.*, 34 (A 13): 9.
1968. *Ditrigona regularis*: Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 451.

Distribution: India: Assam, Khasis, Cherrapunji, Sikkim (Wilkinson, 1968). *Elsewhere:* Burma (Wilkinson, 1968); Nepal (Yazaki, 1993).

42. *Ditrigona sericea* (Leech, 1898)

1898. *Teldenia sericea* Leech, *Trans. Ento. Soc. Lond.*, 1898: 363.
1899. *Drepana fulvicosta* Dudgeon, *J. Bomb. Nat. Hist. Soc.*, 12: 652.
1943. *Leucodrepana nivea birmanica* Bryk, *Ark. Zool.*, 34 (A 13): 7.
1968. *Ditrigona sericea*: Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 434.

Distribution: India: NE India (Darjeeling, Sikkim, Manipur) (Wilkinson, 1968). *Elsewhere:* China, Burma, Nepal (Wilkinson, 1972).

43. *Ditrigona spatulata* Wilkinson, 1968

1968. *Ditrigona spatulata* Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 427-428.

Distribution: India: Sikkim (Tonglo) (Wilkinson, 1968). *Elsewhere:* Nepal (Wilkinson, 1972).

44. *Ditrigona triangularia* (Moore, 1868)

1868. *Urapteryx triangularia* Moore, *Proc. Zool. Soc. Lond.*, 1867: 612.
1968. *Ditrigona triangularia*: Wilkinson, *Trans. Zool. Soc. Lond.*, 31: 448.

Distribution: India: Cherrapunji (Swinhoe, 1895); North-East India, Sikkim, Assam (Wilkinson, 1968). *Elsewhere:* China, Taiwan, Burma, Nepal, Tibet (Wilkinson, 1968).

Genus *Drepana* Schrank, 1802Schrank, 1802; *Fauna Boica*, 2(2): 155.Type-species: *Phalaena falcataria* Linnaeus, 1758.**45. *Drepana dentifera* Warren, 1922**1922. *Drepana dentifera* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 464.*Distribution*: India: Sikkim (Warren, 1922). *Elsewhere*: not known.**46. *Drepana dispilata dispilata* Warren, 1922**1922. *Drepana dispilata* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 463.1943. *Drepana x-z-nigrum* Bryk, 1943; *Ark. Zool.*, **34**(A 13): 11.1968. *Drepana dispilata dispilata*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 109.*Distribution*: India: Sikkim, North India (Watson, 1968). *Elsewhere*: North Burma (Watson, 1968); Nepal (Wilkinson, 1972).*Remark*: The other two Chinese subspecies are: *D. dispilata rufata* Watson, 1968 and *D. dispilata grisearipennis* Strand, 1911.**47. *Drepana ochreipennis* Hampson, 1892**1892. *Drepana ochreipennis* Hampson, *Fauna Br. India (Moths)*, **1**: 335.*Distribution*: India: Sikkim (Hampson, 1892). *Elsewhere*: not known.*Remarks*: Although, The Global Lepidoptera names Index (www.nhm.ac.uk) consider this species under *Watsonalla* Minet but the author of genus *Watsonalla* have not dealt with this species, so far (Pers. Comm.).**48. *Drepana pallida pallida* Moore, 1879**1879. *Drepana pallida* Moore, *Descr. Indian lepid. Atkinson*, (1): 841968 *Drepana pallida pallida*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 107*Distribution*: India: NE India, Darjeeling (Watson, 1968); Uttarakhand (Smetacek, 2002); Arunachal Pradesh (Chandra & Sambath, 2013). *Elsewhere*: North Burma (Watson, 1968); Nepal (Wilkinson, 1972).*Remarks*: *Drepana pallida cretacea* Hampson, 1914 (China, Vietnam), *D. pallida flexuosa* Watson, 1968 (China) and *D. pallida nigromaculata* Okano, 1959 (Taiwan) are other known subspecies.**49. *Drepana rufofasciata* Hampson, 1892**1892. *Drepana rufofasciata* Hampson, *Fauna Br. India (Moths)*, **1**: 334.*Distribution*: India: Sikkim, Tonglu (10,000 ft) (Watson, 1968). *Elsewhere*: China (Watson, 1968); Tibet, Nepal (Wilkinson, 1972).Genus *Drapetodes* Guenée, 1857Guenée, 1857; *Hist. nat. Ins., Spec. gén. Lépid.*, **10**: 424.Type-species: *Drapetodes mitaria* Guenée, 1857.**50. *Drapetodes fratercula* Moore, 1887**1887. *Drapetodes fratercula* Moore, *Lepid. Ceylon*, **3**: 541.1922. *Drapetodes fratercula albidior* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 459.1998. *Drapetodes fratercula*: Holloway, *Moths of Borneo*, **8**: 50.*Distribution*: India: Cherrapunji (Swinhoe, 1895) N.E. Himalaya (Holloway, 1998). *Elsewhere*: Sri Lanka, Borneo, Bali, Sulawesi (Holloway, 1998).**51. *Drapetodes matulata* Felder & Rogenhofer, 1874**1874. *Drapetodes matulata* Felder & Rogenhofer, *Reise öst. Fregatte Novara*, p1. 134.1902. *Drapetodes obliquifasciata* Swinhoe, *Trans. Ent. Soc. Lond.*, 1902: 589.1998. *Drapetodes matulata*: Holloway, 1998; *Moths of Borneo*, **8**: 51.*Distribution*: India: Assam, Khasia (Warren, 1922; Holloway, 1998). *Elsewhere*: Java, Peninsular Malaysia, Borneo (Holloway, 1998).**52. *Drapetodes mitaria* Guenée, 1857**1857. *Drapetodes mitaria* Guenée, *Hist. nat. Ins., Spec. gén. Lépid.*, **9**: 424.1923. *Drapetodes mitaria*: Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 459.*Distribution*: India: Sikkim, Assam, (Warren, 1922); Central India (Holloway, 1998). *Elsewhere*: Taiwan, Singapore (Holloway, 1998).Genus *Euphalacra* Warren, 1897*Euphalacra* Warren, 1897; *Novit. Zool.*, **4**: 195.Type-species: *Euphalacra nigradorsata* Warren, 1897.**53. *Euphalacra nigradorsata* Warren, 1897**1897. *Euphalacra nigradorsata* Warren, *Novit. Zool.*, **4**: 196.*Distribution*: India: Khasis (Warren, 1897); NE Himalaya (Holloway, 1998). *Elsewhere*: not known.Genus *Gogana* Walker, [1866]Walker, [1866]; *List Lep. Het. Br. Mus.*, **34**: 1219.Type-species: *Gogana specularis* Walker, 1865.

54. *Gogana carnosa* (Swinhoe, 1895)

1895. *Phalacropsis carnosa* Swinhoe, *Trans. Ent. Soc. Lond.*, **1895** (1): 5.
1998. *Gogana carnosa*: Holloway, *Moths of Borneo*, **8**: 39.

Distribution: India: Khasis, Cherrapunji (Hampson, 1896; Holloway, 1998). *Elsewhere*: not known.

Genus *Hemiphruda* Warren, 1923

Warren, 1923; In Seitz, *Gross-Schmett. Erde* **10**: 473.

Type-species: *Drepana mecasa* Swinhoe, 1894.

55. *Hemiphruda mecasa* (Swinhoe, 1894)

1894. *Drepana mecasa* Swinhoe, 1894; *Ann. Mag. nat. Hist.*, **6**(14): 434.
1923. *Hemiphruda mecasa*: Warren, in Seitz, *Gross-Schmett. Erde* **10**: 473.

Distribution: India: Khasis, Cherrapunji (Swinhoe, 1894; Warren, 1923). *Elsewhere*: not known.

Genus *Leucoblepsis* Warren, 1922

Warren, 1922; In Seitz, *Gross-Schmett. Erde*, **6**: 462.

Type-species: *Problepsidis carneotincta* Warren, 1901.

56. *Leucoblepsis excisa* (Hampson, 1892)

1892. *Drepana excisa* Hampson, *Fauna Br. India (Moths)*, **1**: 338.
1901. *Problepsidis carneotincta* Warren, *Novit. zool.*, **8**: 191.
1998. *Leucoblepsis excisa*: Holloway, *Moths of Borneo*, **8**: 33.

Distribution: India: N.E. Himalaya (Sikkim, Naga Hills) (Holloway, 1998). *Elsewhere*: Taiwan, Peninsular Malaysia, Sumatra, Borneo (Holloway, 1998).

57. *Leucoblepsis fenestraria* (Moore, 1867)

1867. *Drepanodes fenestraria* Moore, *Proc. Zool. Soc. Lond.*, **1867**: 618.
1903. *Leucoblepsis ostia* Swinhoe, *Fasc. Malay Zool.*, **1**: 59.
1998. *Leucoblepsis fenestraria*: Holloway, *Moths of Borneo*, **8**: 33.

Distribution: India: Bengal (Moore, 1867); N.E. Himalaya (Holloway, 1998). *Elsewhere*: Taiwan; Peninsular Malaysia, Borneo (Holloway, 1998).

Genus *Macrocilix* Butler, 1886

Butler, 1886; *Illustr. Lep. Het. Br. Mus.*, **6**: 18.

Type-species: *Argyris mysticata* Walker, [1863].

58. *Macrocilix mysticata mysticata* (Walker, 1863)

1863. *Argyris mysticata* Walker, *List Lep. Het. Br. Mus.*, **26**: 1617.
1943. *Macrocilix mysticata bidentata* Bryk, *Ark. Zool.*, **34**(A 13): 5.
1958. *Macrocilix mysticata mysticata*: Inoue, *Trans. Shikoku Ent. Soc.*, **6**: 12.

Distribution: India: Dharamshala, Sikkim (Hampson, 1892); North India (Watson, 1868); Arunachal Pradesh

(Chandra & Sambath, 2013); Uttarakhand (Sondhi & Sondhi, 2016). *Elsewhere*: Burma (Watson, 1968); Nepal (Wilkinson, 1972).

Remark: Further, *M. mysticata watsoni* Inoue, 1958 is known from China and Japan whereas *M. mysticata brevinotata* Watson, 1968 is known from China (Watson, 1968).

59. *Macrocilix maia* (Leech, 1888)

1888. *Argyris maia* Leech, *Proc. Zool. Soc. Lond.*, **1888**: 647.
1962. *Macrocilix maia*: Inoue, *Insect Japonica* **2**: 10.

Distribution: India: (Watson, 1968). *Elsewhere*: Korea, Japan, China, Taiwan; Peninsular Malaysia, Sumatra, Borneo (Holloway, 1998).

Genus *Macrauzata* Butler, 1889

Butler, 1889; *Illustr. Lep. Het. Br. Mus.*, **7**: 43.

Type-species: *Comibaena fenestraria* Moore, [1868].

60. *Macrauzata fenestraria* (Moore, 1868)

1868. *Comibaena fenestraria* Moore, *Proc. Zool. Soc. Lond.* **1867**: 639.
1889. *Macrauzata fenestraria*: Butler, *Ill. typical Spec. Lep. Het. Colln Br. Mus.*, **7**: 43.

Distribution: India: Kangra, Sikkim (Hampson, 1892), N. India (Holloway 1998); Uttarakhand (Smetacek, 2002); Arunachal Pradesh (Chandra & Sambath, 2013). *Elsewhere*: Japan (Hampson, 1892); Nepal (Wilkinson, 1972).

Genus *Microblepsis* Warren, 1922

Warren, 1922; *Gross-Schmett. Erde.*, **10**: 461.

Type-species: *Problepsidis cupreogrisea* Hampson, 1895.

61. *Microblepsis prunicolor* (Moore, 1888)

1888. *Drepana prunicolor* Moore, *Descr. Indian lep. Atkinson*, (3): 288.
1943. *Nordstroemia prunicolor warreni* Bryk, *Ark. Zool.*, **34**(A 13): 14.
1968. *Microblepsis prunicolor*: Watson, 1968; *Bull. Br. Mus. nat. Hist. Supp.*, **12**: 53.

Distribution: India: N.E. India (Khasis), Sikkim (Watson 1968). *Elsewhere*: N.E. Burma and China (Watson 1968; Holloway, 1998); Nepal (Wilkinson, 1972)

62. *Microblepsis leucosticta* (Hampson, 1895)

1895. *Drepana leucosticta* Hampson, *Trans. Ent. Soc. Lond.*, **1895**: 287.
1998. *Microblepsis leucosticta*: Holloway, *Moths of Borneo*, **8**: 59.

Distribution: India: NE India, Sikkim, Khasis (Watson, 1968). *Elsewhere*: S. China, Peninsular Malaysia, Sumatra, Borneo (Holloway, 1998).

63. *Microblepsis cupreogrisea* (Hampson, 1895)

1895. *Problepsidis cupreogrisea* Hampson, *Trans. Ent. Soc. Lond.*, 1895: 287.

1998. *Microblepsis cupreogrisea*: Holloway, *Moths of Borneo*, **8**: 59.

Distribution: India: N.E. Himalaya (Holloway, 1998). *Elsewhere*: S. Burma, Borneo, Sumatra (Holloway, 1998).

64. *Microblepsis rugosa* (Watson, 1968)

1968. *Betalbara rugosa* Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 62.

1998. *Microblepsis rugosa*: Holloway, *Moths of Borneo*, **8**: 58.

Distribution: India: NE India, Khasis, Naga Hills (Watson, 1968). *Elsewhere*: Malaysia (Watson, 1968).

Remark: Genus *Microblepsis* was given priority over genus *Betalbara* as mentioned in Holloway (1998)

65. *Microblepsis violacea* (Butler, 1889)

1889. *Agnidra violacea* Butler, *Illustr. Lep. Het. Br. Mus.*, **7**: 42.

1959. *Albara takasago* Okano, *Ann. Rep. Gakugei Fac. Iwate Univ.*, **14** (2): 38. (Syn. By Watson, 1968)

1970. *Microblepsis violacea*: Inoue, *Tinea*, **8**(1): 187.

Distribution: India: N.W. and N.E. India (Dharamshala, Mussorie, Dalhousie, Darjeeling) (Watson, 1968), Uttarakhand (Smetacek, 2002). *Elsewhere*: Taiwan, China (Watson, 1968); Nepal (Wilkinson, 1972).

Genus *Monoprasta* Warren, 1923

Warren, 1923; in Seitz, *Gross-Schmett. Erde*, **10**: 474.

Type-species: *Phalacra nudobia* Swinhoe, 1894.

66. *Monoprasta nudobia* (Swinhoe, 1894)

1894. *Phalacra nudobia* Swinhoe, *Ann. Mag. nat. Hist.*, **6**(14): 435.

1923. *Monoprasta nudobia*: Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 474.

Distribution: India: Khasi Hills, Cherrapunji (Hampson, 1896; Warren, 1923). *Elsewhere*: not known.

Genus *Nordstromia* Bryk, 1943

Bryk, 1943; *Ark. Zool.*, **34** (A 13): 13.

Type-species: *Nordströmia amabilis* Bryk, 1943.

67. *Nordstromia vira* (Moore, 1866)

1866. *Drepana vira* Moore, *Proc. Zool. Soc. Lond.*, **1865** (3): 817.

1894. *Albara erpina* Swinhoe, *Ann. Mag. Nat. Hist.*, **14**(6): 433.

1897. *Albara gracillima* Warren, *Novit. Zool.*, **4**: 12.

1916. *Drepana ocellata* Oberthür, *Étud. Lépid. Comp.*, **12**(1): 375.

1922. *Albara mimetica* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 470.

1943. *Nordströmia amabilis* Bryk, *Ark. Zool.*, **34**(A 13): 13.

1943. *Nordströmia mimetica pallidina* Bryk, *Ark. Zool.*, **34**(A 13): 14.

1968. *Nordstromia vira*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 73.

Distribution: India: Darjeeling, Khasis, Jaintia hills, Nainital, Sikkim (Watson, 1968). *Elsewhere*: N.E. Burma and China (Watson, 1968); Nepal (Wilkinson, 1972).

Remark: Except *A. erpina* (Syn. by Hampson, 1896) all other synonyms by Watson (1968).

68. *Nordstromia bicostata bicostata* (Hampson, 1912)

1912. *Drepana bicostata* Hampson, *J. Bombay nat. Hist. Soc.*, **21**(4): 1272.

1968. *Nordstromia bicostata bicostata*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 75.

Distribution: India: North India, Sikkim (Watson, 1968). *Elsewhere*: N. Burma (Watson, 1968; Wilkinson, 1972).

Remark: *Nordstromia bicostata opalescens* (Oberthur, 1916) known from China.

69. *Nordstromia argenteiceps* (Warren, 1922)

1922. *Albara argenteiceps* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 470.

1968. *Nordstromia argenteiceps*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 77.

Distribution: India: Sikkim and NE India (Khasis) (Watson, 1968). *Elsewhere*: Nepal (Wilkinson, 1972).

70. *Nordstromia lilacina* (Moore, 1888)

1888. *Drepana lilacina* Moore, *Proc. Zool. Soc. Lond.*, 1888: 401.

1968. *Nordstromia lilacina*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 87.

Distribution: India: NE India, NW India (Dharmshala) (Watson, 1968); Uttarakhand (Sondhi & Sondhi, 2016). *Elsewhere*: Nepal (Wilkinson, 1972), Thailand (Buchsbaum, 2010)

71. *Nordstromia simillima* (Moore, 1888)

1888. *Drepana simillima* Moore, *Proc. zool. Soc. Lond.*, 1888: 402.

1968. *Nordstromia simillima*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 87.

Distribution: India: NW India, Dharmshala (Watson, 1968). *Elsewhere*: not known.

72. *Nordstromia duplicata* (Warren, 1922)

1922. *Albara duplicata* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 469.

1922. *Albara olivacea* Warren, in Seitz, *DGross-Schmett. Erde*, **10**: 469.
 1968. *Nordstromia duplicata*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 88.

Distribution: India: N.E. Himalaya (Khasis) (Watson, 1968). *Elsewhere*: W. China, Borneo (Holloway, 1998).

73. *Nordstromia humerata* (Warren, 1896)

1896. *Albara humerata* Warren, *Novit. Zool.* **3**: 335.
 1968. *Nordstromia humerata*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 90.

Distribution: India: NE India, Khasis (Watson, 1968). *Elsewhere*: not known.

Genus *Paralbara* Watson, 1968

Watson, 1968; *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 16.

Type-species: *Fascellina muscularia* Walker, 1866.

74. *Paralbara muscularia* (Walker, 1866)

1866. *Fascellina muscularia* Walker, *List Lep. Het. Br. Mus.*, **35**: 1554.
 1892. *Drepana orphnina* Hampson, *Fauna Br. India (Moths)*, **1**: 337.
 1922. *Albara inaequidiscata* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 469.
 1968. *Paralbara muscularia*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 20.

Distribution: India: Naga Hills, Khasis (Watson, 1968). *Elsewhere*: North Burma, Bhutan, China (Watson, 1968).

75. *Paralbara perhamata* (Hampson, 1892)

1892. *Drepana perhamata* Hampson, *Fauna Br. India (Moths)*, **1**: 336.
 1968. *Paralbara perhamata*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 22.

Distribution: India: NE Himalaya (Jaintia, Khasis, Nagas, Sikkim) (Hampson, 1892; Watson, 1968; Holloway, 1998). *Elsewhere*: Borneo (Holloway, 1998).

Remark: First reference wrongly cited as *Albara* by Holloway (1998).

Genus *Phalacra* Walker, 1866

Walker, 1866; *List Spec. Lepid. Insects Colln Br. Mus.*, **35**: 1638.

Type-species: *Phalacra metagonaria* Walker, 1866.

76. *Phalacra acutipennis* Swinhoe, 1903

1903. *Phalacra acutipennis* Swinhoe, *Ann. Mag. Nat. Hist.*, **11**(7) (65): 502.

Distribution: India: Khasi hills (Swinhoe, 1903; Warren, 1922). *Elsewhere*: not known.

77. *Phalacra albilinea* Warren, 1899

1899. *Phalacra albilinea* Warren, *Novit. Zool.*, **6**: 4.

Distribution: India: Khasi Hills (Warren, 1899; Warren, 1922). *Elsewhere*: not known.

78. *Phalacra excisa* Hampson, 1892

1892. *Phalacra excisa* Hampson, *Fauna Br. India (Moths)*, **1**: 345-346.

Distribution: India: Sikkim (Hampson, 1892; Warren, 1922). *Elsewhere*: not known.

79. *Phalacra ochrea* Warren, 1922

1922. *Phalacra ochrea* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 463.

Distribution: India: Khasi Hills (Warren, 1922). *Elsewhere*: not known.

80. *Phalacra perspicaria* Fabricius, 1798

1798. *Phalaena perspicaria* Fabricius, *Ent. Syst. Suppl.*, **1798**: 449.

Distribution: India: Tranquebar (Coromandel Coast) Tamil Nadu (Fabricius, 1798) *Elsewhere*: not known.

Remark: No other information on the species has been found except, the global Lepidoptera network (Lepindex) which consider it under *Phalacra*.

81. *Phalacra strigata* Warren, 1896

1896. *Phalacra strigata* Warren, *Novit. Zool.*, **3**: 338.
 1897. *Phalacra multilineata* Warren, *Novit. Zool.*, **4**: 16.
 1968. *Phalacra strigata*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.) Suppl.*, **12**: 144.

Distribution: India: North India (Khasis, Cherrapunji) (Watson, 1968). *Elsewhere*: China (Watson, 1968).

82. *Phalacra vidhisara* (Walker, 1860)

1860. *Hemerophila vidhisara* Walker, *List Lep. Het. Br. Mus.*, **21**: 319.
 1866. *Phalacra metagonaria* Walker, *List Lep. Het. Br. Mus.*, **35**: 1639.
 1968. *Phalacra vidhisara*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 143.

Distribution: India: Bombay, Kolkata, Nilgiris (Hampson, 1892; Warren, 1922; Watson, 1968). *Elsewhere*: Sri Lanka (Hampson, 1892).

Genus *Problepsidis* Hampson, 1895

Hampson, 1895, *Trans. Ent. Soc. Lond.*, **1895**: 287.

Type-species: *Problepsidis argyrialis* Hampson, 1895.

83. *Problepsidis argyrialis* Hampson, 1895

1895. *Problepsidis argyrialis* Hampson, *Trans. Ent. Soc. Lond.*, **1895**:

287.

Distribution: India: Naga Hills (Hampson, 1895; Warren, 1922). *Elsewhere:* not known.

Remarks: The species name wrongly mentioned as *argyriatis* instead of *argyrialis* by Warren (1922).

Genus ***Strepsigonia*** (Warren, 1897)

Warren, 1897; *Novit. Zool.*, 4: 17

Type-species: *Strepsigonia nigrimacula* Warren, 1897.

84. ***Strepsigonia diluta*** Warren, 1897

1897. *Tridrepana diluta* Warren, *Novit. Zool.*, 4: 18.

1897. *Tridrepana subobliqua* Warren, *Novit. Zool.*, 4: 18.

1927. *Callidrepana takumukui* Matsumura, *J. Coll. Agric. Hokkaido Imp. Univ.*, 19: 45.

1897. *Strepsigonia diluta fujiensis* Chu & Wang, *Sinozoologica*, 5: 113.

1988. *Strepsigonia diluta takamukui* Matsumura; Inoue, *Tinea*, 12: 126.

1998. *Strepsigonia diluta*: Holloway, *Moths of Borneo*, 8: 29.

Distribution: India: Sikkim, N.E. Himalaya (Holloway, 1998). *Elsewhere:* China, Taiwan, Java, Sumatra (ZSM), Peninsular Malaysia, Borneo, Seram (Holloway, 1998).

85. ***Strepsigonia quadripunctata*** (Walker, 1862)

1862. *Drepana quadripunctata* Walker, *J. Linn. Soc. Lond. (Zool.)*, 6: 175.

1998. *Strepsigonia quadripunctata*: Holloway, *Moths of Borneo*, 8: 30.

Distribution: India: N.E. Himalaya (Naga hills) (Holloway, 1998). *Elsewhere:* Sundaland, Sulawesi, S. Moluccas (Holloway, 1998).

Genus ***Streptoperas*** Hampson, 1895

Hampson, 1895; *Trans. Ent. Soc. Lond.*, 2: 289.

Type-species: *Streptoperas luteata* Hampson, 1895

86. ***Streptoperas luteata*** Hampson, 1895

1895. *Streptoperas luteata* Hampson, *Trans. Ent. Soc. Lond.*, 1895: 289.

1896. *Streptoperas luteatum*: Hampson, *Fauna Br. India, (Moths)*, 4: 478.

Distribution: India: NE Bengal (Hampson, 1895), N.E. Himalaya (Holloway, 1998). *Elsewhere:* Borneo, Sumatra, Java, Bali, Sulawesi (Holloway, 1998).

Genus ***Sewa*** Swinhoe, 1900

Swinhoe, 1900; *Cat. Het. Mus. Oxford*, 2: 591.

Type-species: *Abraxas orbiferata* Walker, 1862.

87. ***Sewa orbiferata*** (Walker, 1862)

1862. *Abraxas orbiferata* Walker, *List Lep. Het. Br. Mus.*, 24: 1126.

1868. *Argyris insignata* Moore, *Proc. zool. Soc. Lond.*, 1867: 645.

1889. *Platypteryx cilicoides* Snellen, *Tijdschr. Ent.*, 32: 9.

1998. *Sewa orbiferata*: Holloway, *Moths of Borneo*, 8: 26.

Distribution: India: Sikkim, Khasis (Hampson, 1892); N.E. Himalaya (Holloway, 1998). *Elsewhere:* Bhutan (Hampson, 1892), Borneo, Peninsular Malaysia, Sumatra (ZSM), Burma, S. & W. China, Java (Holloway, 1998).

Genus ***Teldenia*** Moore, [1883]

Moore, [1883]; *Lepid. Ceylon*, 2(2): 119.

Type-species: *Teldenia alba* Moore, [1883].

88. ***Teldenia specca*** Wilkinson, 1967

1967. *Teldenia specca* Wilkinson, *Trans. R. Ent. Soc. Lond.*, 119(11): 314-315.

Distribution: India: Assam, Khasis, Cherrapunji, (Wilkinson, 1967), N.E. Himalaya (Holloway, 1998). *Elsewhere:* Borneo, Singapore, China, Borneo, Malaysia, Australia, New Britain (Wilkinson, 1967), W. China to Borneo, Sumatra, Palawan, Buru and New Guinea (Holloway, 1998).

89. ***Teldenia vestigiata*** (Butler, 1880)

1880. *Corycia vestigiata* Butler, *Ann. Mag. Nat. Hist.*, 5(6): 222.

1886. *Teldenia vestigiata*: Butler, *Illustr. Lep. Het. Br. Mus.*, 6: 77.

Distribution: India: Shillong, Cherrapunji (Swinhoe, 1985), Darjeeling, Sikkim, Travancore, Karnataka (Wilkinson, 1967); Uttarakhand (Smetacek, 2002). *Elsewhere:* Nepal (Wilkinson, 1972)

Genus ***Thymistadopsis*** Warren, in Seitz, 1922

Warren, 1922; *Gross-Schmett. Erde*, 10: 461.

Type-species: *Problepsidis albidescens* Hampson, 1895.

90. ***Thymistadopsis albidescens*** (Hampson, 1895)

1895. *Problepsidis albidescens* Hampson, *Trans. Ent. Soc. Lond.*, 1895 (2): 288.

1968. *Thymistadopsis albidescens*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.) Suppl.*, 12: 124.

Distribution: India: NE India, Sikkim, Assam, Khasi Hills (Watson, 1968). *Elsewhere:* not known.

91. ***Thymistadopsis trilinearia trilinearia*** (Moore, 1868)

1868. *Drepanodes trilinearia* Moore, *Proc. Zool. Soc. Lond.*, 1867: 618.

1896. *Tridrepana trisulcata* Warren, *Novit. Zool.*, 3: 340.

1968. *Thymistadopsis trilinearia trilinearia*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.) Suppl.*, 12: 126.

Distribution: India: NE India, Sikkim, Darjeeling (Watson, 1968). *Elsewhere:* Nepal (Wilkinson, 1972)

Remark: Further, *Thymistadopsis trilineararia pulvis* (Oberthür, 1916) is known from China.

Genus *Thymistida* Walker, 1865

Walker, 1865; *List Lep. Het. Br. Mus.*, **32**: 515.

Type-species: *Thymistida tripunctata* Walker, 1865.

92. *Thymistida nigritincta* Warren, 1923

1923. *Thymistida nigritincta* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 474.

1923. *Thymistida rufa* Warren, in Seitz, *ibid.*, **10**: 475.

1943. *Hybodrepana grotesca* Bryk, *Ark. Zool.*, **34A**, (13): 23.

1968. *Thymistida nigritincta*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.) Suppl.*, **12**: 27.

Distribution: India: NE India, Khasis (Watson, 1968). *Elsewhere:* N. Burma (Watson, 1968).

93. *Thymistida tripunctata* Walker, 1865

1865. *Thymistida tripunctata* Walker, *List Lep. Het. Br. Mus.*, **32**: 515.

1867. *Erosia cervinaria* Moore, *Proc. Zoo. Soc. Lond.*, **1867**: 646.

1943. *Thymistida nigritincta divisa* Bryk, *Ark. Zool.*, **34 A**, (13): 23.

1968. *Thymistida tripunctata*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.) Suppl.*, **12**: 27.

Distribution: India: Sikkim (Hampson, 1892), NE India, Arunachal Pradesh (Chandra & Sambath, 2013). *Elsewhere:* Burma, China (Watson, 1968), Nepal (Wilkinson, 1972; Yazaki, 1992).

94. *Thymistida undilineata* Warren, 1923

1923. *Thymistida undilineata* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 474.

Distribution: India: NE India, Khasis (Warren, 1923; Watson, 1968). *Elsewhere:* not known.

Genus *Tridrepana* Swinhoe, 1895

Swinhoe, 1895, *Trans. Ent. Soc. Lond.*, **1895**: 3.

Type-species: *Drepana albonotata* Moore, 1879.

95. *Tridrepana acuta* Watson, 1957

1957. *Tridrepana acuta* Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**, 430-431.

Distribution: India: South India (but needs confirmation) (Watson, 1957). *Elsewhere:* Sri Lanka (Watson, 1957).

96. *Tridrepana adelpha* Swinhoe, 1905

1905. *Tridrepana adelpha* Swinhoe, *Ann. Mag. nat. Hist.*, **16**(7): 620.

Distribution: India: North East India, Cherrapunji, Khasis (Watson, 1957). *Elsewhere:* Nepal (Wilkinson, 1972).

97. *Tridrepana albonotata albonotata* (Moore, 1879)

1879. *Drepana albonotata* Moore, *Descr. Indian lepid. Atkinson*, **1**: 83.

1886. *Callidrepana ochrea* Butler, *Illustr. Lep. Het. Br. Mus.*, **6**: 17.

1922. *Iridrepana glaciata* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 467.

1957. *Tridrepana albonotata albonotata*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.)*, **4**: 436.

Distribution: India: South India, Sikkim, Darjeeling (Watson, 1957), Uttarakhand (Smetacek, 2002). *Elsewhere:* Vietnam (Watson, 1957).

Remarks: Apart from this and below mentioned species, there are four other subspecies: *T. albonotata angusta* Watson, 1957 (Borneo, Sumatra, Malaysia); *T. albonotata rotunda* Watson, 1957 (Bali, Java); *T. albonotata celebesensis* Watson, 1957 (Sulawesi); *T. albonotata ferrea* (Hampson, 1892) Sri Lanka. However, Holloway (1998) considered all the species under *T. albonotata* without mentioning *Syn. Nov.*, though in distribution he mentioned different subspecies restricted to specific areas.

98. *Tridrepana albonotata pervasata* (Warren, 1922)

1922. *Iridrepana septempunctata pervasata* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 465.

1957. *Tridrepana albonotata pervasata*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**: 444.

Distribution: India: Travancore, South India (Watson, 1957). *Elsewhere:* not known.

99. *Tridrepana flava* (Moore, 1879)

1879. *Tridrepana flava* Moore, *Descr. Indian lepid. Atkinson*, **1**: 84.

1957. *Tridrepana flava contracta* Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**: 495-496.

1957. *Tridrepana flava unita* Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**: 496-497.

1998. *Tridrepana flava*: Holloway, *Moths of Borneo*, **8**: 55.

Distribution: India: Sikkim, Cherrapunji, Khasis, Darjeeling (Watson, 1957), N.E. Himalaya (Holloway, 1998). *Elsewhere:* Taiwan, Sundaland, Sulawesi (Holloway, 1998).

100. *Tridrepana fulva* (Hampson, 1892)

1892. *Drepana fulva* Hampson, *Fauna Br. India (Moths)*, **1**: 342.

1957. *Tridrepana fulva*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**: 489.

Distribution: India: Sikkim (Yatong, 13000 ft.) (Hampson, 1892). *Elsewhere:* Nepal, Tibet, China (Watson, 1957; Wilkinson, 1972).

101. *Tridrepana fulvata brevis* (Watson, 1957)

1957. *Tridrepana fulvata brevis* Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**: 422.

Distribution: India: Assam, Khasis (Watson, 1957). *Elsewhere*: China, Myanmar (Song *et al.*, 2011).

Remark: The nominate subspecies is known from Sumatra, Java, Bali, Malaysia, Borneo.

102. *Tridrepana postica* (Moore, 1879)

1879. *Drepana postica* Moore, *Lepid. Atk.* **1**: 84.
1892. *Drepana xanthoptera* Hampson, *Fauna Br. India (Moths)*, **1**: 341.
1957. *Tridrepana postica*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**: 472.

Distribution: India: North East India (Cherrapunji, Shillong, Khasi Hills, Naga hills) (Watson, 1957). *Elsewhere*: Malaysia (Watson, 1957).

103. *Tridrepana rubromarginata indica* Watson, 1957

1957. *Tridrepana rubromarginata indica* Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **4**: 486.

Distribution: India: Sikkim, Tonglu (10,000ft) (Watson, 1957). *Elsewhere*: Bhutan, Nepal (Watson, 1957); Tibet (Song *et al.*, 2011).

Remark: The nominate subspecies is known from China.

104. *Tridrepana sadana* (Moore, 1865)

1865. *Drepana sadana* Moore, *Proc. Zool. Soc. Lond.*, **1865**: 817.
1943. *Tridrepana adeipha matronalis* Bryk, *Ark. Zool.*, **34** A, (13): 16.
1957. *Tridrepana sadana*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.)*, **4**: 482.

Distribution: India: North India (Kumaon), Sikkim, Shillong (Watson, 1957). *Elsewhere*: North East Burma (Watson, 1957); Nepal (Wilkinson, 1972).

105. *Tridrepana septempunctata septempunctata* Warren, 1896

1896. *Tridrepana septempunctata* Warren, *Novit. Zool.* **3**: 339.
1957. *Tridrepana septempunctata septempunctata*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.)*, **4**: 458.

Distribution: India: Assam (Jaintia Hills) (Watson, 1957). *Elsewhere*: not known.

Remark: *Tridrepana septempunctata nitidior* Watson, 1957 is known from Sumatra.

Genus *Zusidava* Walker, [1863]

Walker, [1863]; *List Spec. Lepid. Insects Colln Br. Mus.*, **26**: 1637.

Type-species: *Zusidava tortricaria* Walker, [1863].

106. *Zusidava sinuosa* (Moore, 1888)

1888. *Emodesa sinuosa* Moore, *Descr. Indian lep. Atkinson*, (3): 256.
1998. *Zusidava sinuosa*: Holloway, *Moths of Borneo*, **8**: 22.

Distribution: India: N.E. Himalaya (Khasi Hills) (Moore, 1888). *Elsewhere*: Burma (Holloway, 1998).

Subfamily: ORETINAE

Genus *Neoreta* Warren, 1897

Neoreta Warren, 1897, *Novit. Zool.*, **4**: 378.

Type-species: *Oreta olga* Swinhoe, 1894.

107. *Neoreta castanea* (Hampson, 1891)

1891. *Oreta castanea* Hampson, *Illustr. Lep. Het. Br. Mus.*, **8**: 9.
1891. *Oreta rotundipex* Hampson, *Illustr. Lep. Het. Br. Mus.*, **8**: 9. (Syn. by Watson, 1967).
2012. *Neoreta castanea*: Song *et al.*, *Zootaxa*, **3445**: 30.

Distribution: India: South India, Nilgiris (Watson, 1967). *Elsewhere*: not known.

108. *Neoreta olga* (Swinhoe, 1894)

1894. *Oretaolga* Swinhoe, *Ann. Mag. nat. Hist.*, (6) **14**(84): 434.
2012. *Neoreta olga*: Song *et al.*, *Zootaxa*, **3445**: 30.

Distribution: India: North East India, Shillong, Khasis (Watson, 1967). *Elsewhere*: China (Song *et al.*, 2012).

Genus *Oreta* Walker, 1855

Oreta Walker, 1855, *List Lep. Het. Br. Mus.*, **5**: 1166.

Type-species: *Oreta extensa* Walker, 1855.

109. *Oreta extensa* Walker, 1855

1855. *Oreta extensa* Walker, *List Lep. Het. Br. Mus.*, **5**: 1166.
1905. *Oreta figlina* Swinhoe, *Ann. Mag. Nat. Hist.*, (7) **16** (92): 142.
1967. *Oreta extensa*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **19**(3): 200.

Distribution: India: NE India (Cherrapunji, Shillong), South India (Watson, 1961, 1967). *Elsewhere*: Sumatra, Java, Sri Lanka, Taiwan, Sulawesi (Watson, 1967); China, Tibet, Japan, Thailand, Indonesia (Song *et al.*, 2012).

110. *Oreta griseotincta griseotincta* Hampson, 1892

1892. *Oreta griseotincta* Hampson, *Fauna Br. India (Moths)*, **1**: 350.
1899. *Oreta olivacea* Dudgeon, *J. Bombay nat. Hist. Soc.*, **12**(4): 657.
1923. *Oreta carnea nucicolor* Warren, in Seitz, *Gross-Schmett. Erde*, **10**: 484.
1927. *Mimoreta horishana* Matsumura, *J. Coll. Agric. Hokkaido Imp. Univ.*, **19**(1): 46.
1967. *Oreta griseotincta griseotincta*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, **19**(3): 203.
1988. *Oreta changi* Inoue, *Tinea*, **12**(15): 133 (Song *et al.*, 2012)

Distribution: India: NE India, Sikkim, Khasis (Watson, 1961, 1967). *Elsewhere:* Taiwan (Watson, 1967), China, Hongkong, India, Malaysia, Singapore (Song *et al.*, 2012).

Remark: *Oreta griseotincta acutior* Watson, 1961 is known from Malaysia and Singapore.

111. *Oreta obtusa obtusa* (Walker, 1855)

1855. *Oreta obtusa* Walker, *List Lep. Het. Br. Mus.*, 5: 1167.

1892. *Oreta obliquilinea* Hampson, *Fauna Brit. India, Moths*, 1: 349.

1923. *Psiloreta inconspicua* Warren, in Seitz, *Gross-Schmett. Erde*, 10: 486.

1961. *Psiloreta obtusa obtusa*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, 10: 345.

Distribution: India: Kashmir, Sikkim, Naga hills, Punjab (Hampson, 1892), North India, Mussoorie (Watson, 1961; 1967). *Elsewhere:* Burma, China, Nepal, Indonesia (Watson, 1961, 1967; Song *et al.*, 2012).

Remarks: Other known subspecies not reported from India are: *N.o. aequitermen* Watson, 1961 (Malaya, Sumatra, Celebes); *N.o. javae* Watson, 1961 (Java, Bali); *N.o. dejeani* Watson, 1967 and its synonyms *Oreta sinuata* Chu & Wang, 1987 and *Oreta asignis* Chu & Wang, 1987 (China, Tibet). Further, *N.o. speciosa* Bryk, 1943 (Mynamar, NE Burma, China, Tibet) raised to species status by Song *et al.* (2012).

112. *Oreta pavaca pavaca* Moore, 1866

1866. *Oreta pavaca* Moore, *Proc. Zool. Soc. Lond.*, 1865 (3): 815.

1923. *Psiloreta pavaca purpurea* Warren, in Seitz, *Gross-Schmett. Erde*, 10: 486.

1923. *Psiloreta pavaca olivacea* Warren, in Seitz, *Gross-Schmett. Erde*, 10: 486.

1967. *Oreta pavaca pavaca*: Watson, *Bull. Br. Mus. nat. Hist. (Ent.)*, 19(3): 182.

1987. *Oreta zigzaga* Chu & Wang, *Acta Ent. Sin.*, 30(3): 296. (Syn. by Song *et al.*, 2012)

Distribution: India: NW India (Shimla, Mussorie), North India, Sikkim, Darjeeling, Khasis (Watson, 1967). *Elsewhere:* China, Tibet, Nepal (Wilkinson, 1972; Song *et al.*, 2012).

Remark: Another known subspecies, *O. pavaca sinensis* Watson, 1967 also includes *O. fusca* Chu & Wang, 1987, *O. unichroma* Chu & Wang, 1987 and *O. lushansis* Fang, 2003 as its synonyms is known from China (Song *et al.*, 2012).

113. *Oreta sanguinea* Moore, 1879

1879. *Oreta sanguinea* Moore, *Descr. Indian lep. Atkinson*, (1): 85.

Distribution: India: N.E. India, Sikkim, Khasis, Darjeeling (Watson, 1967; Song *et al.*, 2012); Arunachal Pradesh (Chandra & Sambath, 2013). *Elsewhere:* Nepal, Tibet (Wilkinson, 1972).

114. *Oreta suffusa* Walker, 1855

1855. *Oreta suffusa* Walker, *List Lep. Het. Br. Mus.*, 5: 1167.

1891. *Oreta violacea* Hampson, *Illustr. Lep. Het. Br. Mus.*, 8: 9.

1967. *Oreta suffusa*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.)*, 19(3): 200-201.

Distribution: India: South India, Nilgiris (Watson, 1967). *Elsewhere:* Sri Lanka (Watson, 1967).

115. *Oreta vatama vatama* (Moore, 1866)

1866. *Oreta vatama* Moore, *Proc. Zool. Soc. Lond.*, (3): 816.

1967. *Oreta vatama vatama*: Watson, *Bull. Br. Mus. Nat. Hist. (Ent.)*, 19 (3): 189.

Distribution: India: NE India, Darjeeling, Sikkim (Watson, 1967). *Elsewhere:* Bhutan, N. Burma (Watson, 1967); Nepal (Wilkinson, 1972).

Remark: Apart from below mentioned subspecies, two other subspecies are: *O. vatama acutula* Watson, 1967 (China, Tibet) and *O. vatama tsina* Watson, 1967 (China).

116. *Oreta vatama luculenta* Watson, 1967

1967. *Oreta vatama luculenta* Watson, *Bull. Br. Mus. Nat. Hist. (Ent.)*, 19(3): 189.

Distribution: India: NW. India (Dalhousie, Shimla), Kashmir (Liddar valley, Gulmarg) (Watson, 1967). *Elsewhere:* Pakistan (Watson, 1967).

Genus *Spectroreta* Warren, 1903

Spectroreta Warren, 1903, *Novit. Zool.*, 10: 255.

Type-species: *Oreta hyalodisca* Hampson, 1896.

117. *Spectroreta hyalodisca* (Hampson, 1896)

1896. *Oreta hyalodisca* Hampson, *Fauna Br. India, (Moths)*, 4: 479.

1903. *Spectroreta hyalodisca*: Warren, *Novit. Zool.*, 10: 255.

1987. *Spectroreta fenestra* Chu & Wang, *Acta Ent. Sin.*, 31(3): 291. (Syn. by Song *et al.* 2012).

Distribution: India: N.E. Himalaya, Khasis (Watson, 1967; Holloway, 1998). *Elsewhere:* Sri Lanka, S. China, Burma, Peninsular Malaysia, Sumatra, Borneo, Kei Is. (Holloway, 1998); China (Song *et al.*, 2012).

Subfamily: THYATIRINAE

Genus *Epipsestis* Matsumura, 1921

Matsumura, 1921; *Thous. Ins. Japan (Additam.)*, 4: 857.

Type-species: *Polyploca ornata* Leech, [1889].

118. *Epipsestis albidisca* (Warren, 1888)

1888. *Polyploca albidisca* Warren, *Proc. Zool. Soc. Lond.*, **1888**: 299.

1888. *Palimpsestis albidisca* Moore, *Proc. Zool. Soc. Lond.*, **1888**: 408.

1896. *Polyploca polychromata* Hampson, *Fauna Br. India (Moths)* **4**: 463.

1982. *Epipsestis albidisca*: Yoshimoto, *Tyo to Ga*, **32** (3,4): 126-128.

Distribution: India: Berham Gully (type locality), Kangra, Dharmshala, Sikkim (László *et al.*, 2007). *Elsewhere*: China (Yunnan, Tibet), Nepal, Pakistan, Vietnam, Thailand (László *et al.*, 2007).

119. *Epipsestis bilineata bilineata* (Warren, 1915)

1915. *Polyploca bilineata* Warren, *Novit. Zool.*, **22**(1): 159.

1915. *Polyploca acutangula* Warren, *Novit. Zool.*, **22** (1): 159.

1984. *Epipsestis bilineata bilineata*: Yoshimoto, *Japan Hetero. J.*, **128**: 33.

Distribution: India: Sikkim, Darjeeling (Warren, 1915). *Elsewhere*: Nepal (László *et al.*, 2007).

Remark: Further subspecies is *E. bilineata pallida* Yoshimoto, 1984 endemic to Taiwan (Yoshimoto, 1984a; László *et al.*, 2007).

120. *Epipsestis castaneata* (Warren, 1915)

1915. *Polyploca castaneata* Warren, *Novit. Zool.*, **22**(1): 157.

2007. *Epipsestis castaneata*: László *et al.*, *Esperiana*, **13**: 299.

Distribution: India: Sikkim (Warren, 1915). *Elsewhere*: China (Hunan), Nepal, Vietnam (László *et al.*, 2007).

121. *Epipsestis dubia* (Warren, 1888)

1888. *Polyploca dubia* Warren, *Proc. Zool. Soc. Lond.*, **1888** (3): 300.

1915. *Polyploca nigrifasciata* Warren, *Novit. zool.*, **22**(1): 158.

1915. *Polyploca nigrifasciata ab. semifascia* Warren, *Novit. Zool.*, **22**(1): 158.

1915. *Polyploca nigrifasciata ab. nigrilineata* Warren, *Novit. Zool.* **22**(1): 158.

2007. *Epipsestis dubia*: Yoshimoto, 1982; *Tyo to Ga*, **32**(3,4): 123-125.

Distribution: India: Sikkim, (Warren, 1888), Kashmir (Warren, 1888; Yoshimoto, 1982; László *et al.*, 2007; Jiang *et al.*, 2015). *Elsewhere*: China, Tibet, Nepal, Pakistan, Myanmar, Vietnam (László *et al.*, 2007).

122. *Epipsestis longipennis* (Yoshimoto, 1982)

1982. *Epipsestis longipennis* Yoshimoto, *Tyo to Ga*, **32**(3-4): 129.

Distribution: India: Sikkim (László *et al.*, 2007) *Elsewhere*: Nepal (László *et al.*, 2007)

123. *Epipsestis nikkoensis* (Matsumura, 1921)

1921. *Polyploca nikkoensis* Matsumura, *Thous. Ins. Japan (Additam.)*, **4**: 856.

1982. *Epipsestis nikkoensis*: Yoshimoto, *Tyo to Ga*, **32**(3,4): 136.

Distribution: India: (László *et al.*, 2007). *Elsewhere*: China, Taiwan, Southeast of Russia, Japan, Korean Peninsula, Nepal, Bhutan, Myanmar, Vietnam, Thailand (László *et al.*, 2007).

Remark: It needs confirmation from India, however its occurrence in the Indian Himalayas is highly likely.

124. *Epipsestis renalis* (Moore, 1888)

1888. *Palimpsestis renalis* Moore, *Proc. Zool. Soc. Lond.*, **1888**: 407.

1982. *Epipsestis renalis*: Yoshimoto, *Tyo to Ga*, **32**(3,4): 136.

Distribution: India: Kangra, Dharamshala (Moore, 1888; Yoshimoto, 1982). *Elsewhere*: China (Hunan), Nepal, Pakistan, Vietnam, Thailand. (László *et al.*, 2007).

125. *Epipsestis witti* László, Ronkay & Ronkay, 2007

2007. *Epipsestis witti* László, Ronkay & Ronkay, *Esperiana*, **13**: 309.

Distribution: India: (Jiang *et al.*, 2015). *Elsewhere*: China (Tibet), Nepal, Vietnam, Thailand (László *et al.*, 2007; Jiang *et al.*, 2015); Myanmar (László *et al.*, 2007).

Remarks: It needs confirmation from India as László *et al.* (2007) do not list specimens from India in the type series of *E. witti*. In addition, Jiang *et al.* (2015) do not specify collecting locality of the species when referring it from India. Nevertheless, the species most probably occurs in certain parts of the Indian Himalayas.

Genus *Euparyphasma* Fletcher, 1979

(replacement name for *Lithocharis* Warren, 1912)

Fletcher, 1979; *Gen. Names Moths World*, **3**: 83.

Type-species: *Polyploca albibasis* Hampson, 1893.

126. *Euparyphasma albibasis albibasis* (Hampson, 1892)

1892. *Polyploca albibasis* Hampson, *Fauna Br. India (Moths)*, **1**: 184.

2007. *Euparyphasma albibasis*: László *et al.*, *Esperiana* **13**: 140.

Distribution: India: Shimla, Sikkim (Hampson, 1892; László *et al.*, 2007). *Elsewhere:* not known.

Remarks: A subspecies *E. albibasis cinereofusca* (Houlbert, 1921) is reported from China and Vietnam only. Sondhi & Sondhi (2016) erroneously cited the author as Prout for *E. albibasis cinereofusca* (Houlbert, 1921). In addition, their report of *E. cinereofusca* is incorrect, as it refers to the nominotypical subspecies.

127. *Euparyphasma maxima* (Leech, 1889)

1889. *Cymatophora maxima* Leech, *Proc. Zool. Soc. Lond.*, **1888**: 653.

2007. *Euparyphasma maxima*: László *et al.*, *Esperiana* **13**: 143.

Distribution: India: Locality not specified (Kim *et al.*, 2006). *Elsewhere:* Korea, Japan, China (László *et al.*, 2007; Kim *et al.*, 2006).

Remark: Kim *et al.* (2006) did not specify its distribution in India, therefore, the occurrence of this species in India requires confirmation.

Genus *Gaurena* Walker, 1865

Walker, 1865; *List Lep. Het. Br. Mus.*, **32**: 619.

Type-species: *Gaurena florens* Walker, 1865.

128. *Gaurena albifasciata* Gaede, 1930

1930. *Gaurena albifasciata* Gaede, In: Seitz, *Gross-Schmetterl.*, **10**: 660.

1966. *Gaurena albifasciata nepalensis* Werny, *Untersuchungen über die Systematik* (Lepidoptera, Thyatiridae): 106.

2007. *Gaurena albifasciata*: László *et al.*, *Esperiana*, **13**: 67.

Distribution: India: (László *et al.*, 2007; Jiang *et al.*, 2015). *Elsewhere:* China (Tibet), Nepal (László *et al.*, 2007).

Remark: Neither László *et al.* (2007), nor Jiang *et al.* (2015) refer to examined specimen from India, therefore the occurrence in the country requires confirmation.

129. *Gaurena argentisparsa* Hampson, 1896

1896. *Gaurena argentisparsa* Hampson, *Fauna Br. India (Moths)*, **4**: 462.

1966. *Gaurena argentisparsa eberti* Werny, *Untersuchungen über die Systematik* (Lepidoptera, Thyatiridae): 114.

2007. *Gaurena argentisparsa*: László *et al.*, *Esperiana*, **13**: 65.

Distribution: India: Sikkim, Yatong (Hampson, 1896). *Elsewhere:* China (Tibet), Nepal, Bhutan (László *et al.*, 2007).

130. *Gaurena aurofasciata* Hampson, 1892

1892. *Gaurena aurofasciata* Hampson, *Fauna Br. India (Moths)*, **1**: 181.

1966. *Gaurena aurofasciata bryki* Werny, *Untersuchungen über die Systematik* (Lepidoptera, Thyatiridae): 96.

2007. *Gaurena aurofasciata*: László *et al.*, *Esperiana* **13**: 72.

Distribution: India: Sikkim (Hampson, 1892; László *et al.*, 2007). *Elsewhere:* China (Yunnan), Myanmar (László *et al.*, 2007).

131. *Gaurena florens* Walker, 1865

1865. *Gaurena florens* Walker, *List Lep. Het. Br. Mus.*, **32**: 620.

1966. *Gaurena florensoliva* Werny, *Untersuchungen über die Systematik* (Lepidoptera, Thyatiridae): 82.

1966. *Gaurena florensobscura* Werny, *ibid.*: 80.

1966. *Gaurena florens yuennanensis* Werny, *ibid.*: 79.

2007. *Gaurena florens*: László *et al.*, *Esperiana*, **13**: 72.

Distribution: India: Darjeeling, Sikkim (Walker, 1865; László *et al.*, 2007); Uttar Pradesh, Himachal Pradesh (László *et al.*, 2007), Arunachal Pradesh (Chandra & Sambath, 2013). *Elsewhere:* China (Tibet, Yunnan, Sichuan), Nepal, Bhutan, Myanmar, Vietnam, Thailand (László *et al.*, 2007; Jiang *et al.*, 2015).

132. *Gaurena florescens* Walker, 1865

1865. *Gaurena florescens* Walker, *List Lep. Het. Br. Mus.*, **32**: 620.

1966. *Gaurena florescens albomaculata* Werny, *Untersuchungen über die Systematik* (Lepidoptera, Thyatiridae): 87.

1966. *Gaurena florescens burmanica* Werny, *ibid.*: 85.

1966. *Gaurena gemella flavescens* Werny, *ibid.*: 90.

2007. *Gaurena florescens*: László *et al.*, *Esperiana*, **13**: 63.

Distribution: India: Sikkim, Khasis, Nagas (Hampson, 1892), Darjeeling (Walker, 1865; László *et al.*, 2007; Jiang *et al.*, 2015). *Elsewhere:* China (Shaanxi, Sichuan, Yunnan, Jiangxi), Nepal, Thailand, Vietnam (László *et al.*, 2007); China (Hubei, Hunan, Guangdong, Sichuan, Yunnan, Tibet), Myanmar, Bangladesh, Cambodia (Jiang *et al.*, 2015).

133. *Gaurena forsteri* Werny, 1966

1966. *Gaurena forsteri* Werny, *Untersuchungen über die Systematik* (Lepidoptera, Thyatiridae): 101.

Distribution: India: Sikkim (László *et al.*, 2007; Jiang *et al.*, 2015).

Elsewhere: Nepal (László *et al.*, 2007); China (Tibet), Nepal (Jiang *et al.*, 2015).

134. *Gaurena grisescens* Oberthür, 1894

1894. *Gaurena grisescens* Oberthür, *Etud. d' Ent.*, **18**: 43.

Distribution: India: Sikkim (Warren, 1922). *Elsewhere:*

China (Sichuan, Yunnan, Tibet) (László *et al.*, 2007; Jiang *et al.*, 2015).

Remarks: Although both László *et al.* (2007) and Jiang *et al.* (2015) restricted the distribution of this species for China only, Warren (1922) mentioned a male specimen collected in Sikkim. Nevertheless, due to the difficult identification of the species of the genus, the occurrence of *G. grisescens* in India needs confirmation.

Genus ***Habrosyne*** Hübner, 1821

Hübner, [1821]; *Verz. bek. Schmett.* (15): 236.

Type-species: *Phalaena derasa* Linnaeus, 1767.

135. *Habrosyne armata* Moore, 1882

1882. *Habrosyne armata* Moore, *Lep. Atk.*, 2: 90.

Distribution: India: Khasi Hills (Moore, 1882; Hampson, 1892); Darjeeling (László *et al.*, 2007). *Elsewhere:* not known.

136. *Habrosyne fraterna* Moore, 1888

1888. *Habrosyne fraterna* Moore, *Proc. Zool. Soc. Lond.*, 1888: 406.

Distribution: India: Kangra, Himachal Pradesh (Moore, 1888). *Elsewhere:* Pakistan (Murree Hills), Nepal (László *et al.*, 2007)

137. *Habrosyne indica indica* (Moore, 1867)

1867. *Gonophora indica* Moore, *Proc. Zool. Soc. Lond.*, 1867: 44.

1943. *Habrosyne fraterna malaisei* Bryk, *Arkiv för Zoologi*, 34 A: 6.

1966. *Habrosyne fraterna chekiangensis* Werny, *Untersuchungen über die Systematik* (*Lepidoptera, Thyatiridae*): 275.

1966. *Habrosyne fraterna japonica* Werny, *ibid.*: 276.

1966. *Habrosyne indica grisea* Werny, *ibid.*: 279.

1966. *Habrosyne indica flavescens* Werny, *ibid.*: 282.

1966. *Habrosyne indica aurata* Werny, *ibid.*: 280.

2007. *Habrosyne indica indica*: László *et al.*, *Esperiana* 13: 193.

Distribution: India: Sikkim (Moore, 1867; László *et al.*, 2007; Jiang *et al.*, 2015). *Elsewhere:* China (Tibet, Yunnan, Shaanxi), Japan, Nepal, Myanmar, Vietnam, Thailand (László *et al.*, 2007; Jiang *et al.*, 2015).

Remarks: Hampson (1892) synonymised *H. indica* (Moore) and *H. fraterna* Moore from India under European and West Palaearctic species *Phalaena Noctua derasa* Linnaeus, 1767 (= *H. (Phalaena) pyritoides* Hufnagel, 1766). However, both former species are obviously valid and thus the latter one is not included in this list. Further subspecies is *H. indica formosana* Werny, 1966 endemic to Taiwan.

138. *Habrosyne intermedia conscripta* Warren, 1912

1912. *Habrosyne conscripta* Warren, in Seitz, *Gross-Schmetterlinge der Erde*, 2: 323.

1966. *Habrosyne conscripta nepalensis* Werny, *Untersuchungen über die Systematik* (*Lepidoptera, Thyatiridae*): 268. (László *et al.*, 2007)

1912. *Habrosyne intermedia conscripta*: László *et al.*, 2007; *Esperiana*, 13: 191.

Distribution: India: Himachal Pradesh (László *et al.*, 2007; Jiang *et al.*, 2015). *Elsewhere:* China, Nepal (László *et al.*, 2007; Jiang *et al.*, 2015).

Remark: The nominate subspecies of *H. intermedia* was described from the Russian Far East (Sikhote Alin) occurring also in northern China, the Korean Peninsula and North Japan (László *et al.*, 2007).

139. *Habrosyne plagiosa* Moore, 1882

1882. *Habrosyne plagiosa* Moore, *Lep. Atk.*, 2: 90.

Distribution: India: Sikkim (Moore, 1882; Hampson, 1892; László *et al.*, 2007). *Elsewhere:* Nepal (Yoshimoto, 1993; László *et al.*, 2007).

140. *Habrosyne sanguinea* Moore, 1882

1882. *Habrosyne sanguinea* Moore, *Lep. Atk.*, 2: 90.

Distribution: India: Sikkim, Darjeeling (Moore, 1882; László *et al.*, 2007; Jiang *et al.*, 2015). *Elsewhere:* China (Tibet), Nepal (László *et al.*, 2007; Jiang *et al.*, 2015).

141. *Habrosyne violacea argenteipuncta* (Hampson, 1892)

1892. *Habrosyne argenteipuncta* Hampson, *Fauna Br. India (Moths)*, 1: 179.

1966. *Habrosyne argenteipuncta nigricans* Werny, *Untersuchungen über die Systematik* (*Lepidoptera, Thyatiridae*): 294.

1966. *Habrosyne argenteipuncta burmanica* Werny, *ibid.*: 297.

2004. *Habrosyne violacea argenteipuncta*: Zhao, *Fauna Sinica, Insecta*, Vol. 36.

Distribution: India: Sikkim (Hampson, 1892; László *et al.*, 2007). *Elsewhere:* China (Yunnan, Tibet) (Jiang *et al.*, 2015). Nepal, Vietnam, Myanmar (László *et al.*, 2007).

Remarks: László *et al.* (2007) considered only two valid subspecies: the nominate subspecies (China, Russian Far East, Korea) and the subspecies *argenteipuncta* Hampson, 1892.

Genus *Horithyatira* Matsumura, 1933

Matsumura, 1933; *Insecta Matsumurana*, 7: 193.

Type-species: *Thyatira decorata* Moore, 1881.

142. *Horithyatira decorata decorata* (Moore, 1881)

1881. *Thyatira decorata* Moore, *Proc. Zool. Soc. Lond.*, **1881**: 328.

1944. *Horithyatira decorata birmanica* Bryk, *Entomologisk Tidskrift*, **65**: 224. (Holloway, 1998)

1966. *Horithyatira decorata thodungensis* Werny, *Untersuchungen über die Systematik* (*Lepidoptera, Thyatiridae*): 140. (Holloway, 1998)

1966. *Horithyatira hoenei* Werny, *Untersuchungen über die Systematik* (*Lepidoptera, Thyatiridae*): 149. (Holloway, 1998)

2007. *Horithyatira decorata decorata*: László *et al.*, *Esperiana*, **13**: 54.

Distribution: India: Darjeeling, Sikkim (Moore, 1881; László *et al.*, 2007). *Elsewhere*: China, Nepal, Bhutan, Myanmar, Vietnam, Thailand (László *et al.*, 2007; Jiang *et al.*, 2015; Zhuang *et al.*, 2017).

Remark: Further valid subspecies are *H. decorata takamukui* (Matsumura, 1921) from Taiwan and *H. decorata kawamurae* (Matsumura, 1921) from Japan.

Genus *Isopsestis* Werny, 1968

Werny, 1968: *Ergebnisse der ForschUnternehmens Nepal Himalaya*, 3: 113.

Type-species: *Palimpsestis cuprina* Moore, 1881.

143. *Isopsestis cuprina* (Moore, 1881)

1881. *Palimpsestis cuprina* Moore, *Proc. Zoo. Soc. Lond.*, **1881**: 331.

2007. *Isopsestis cuprina*: László *et al.*, *Esperiana*, **13**: 168.

Distribution: India: Darjeeling (Moore, 1881; László *et al.*, 2007). *Elsewhere*: China (Tibet), Nepal (László *et al.*, 2007; Jiang *et al.*, 2015).

Genus *Macrothyatira* Marumo, 1916

Marumo, 1916; *Insect World, Gifu, Japan*, **20** (2): 48.

Type-species: *Thyatira flavida* Butler, 1885.

144. *Macrothyatira stramineata stramineata* (Warren, 1912)

1912. *Thyatira stramineata* Warren, In Seitz, *Gross Schmett. Erde*, **2**: 324.

2007. *Macrothyatira stramineata stramineata*: László *et al.*, *Esperiana*, **13**: 79.

Distribution: India: Khasi hills (Warren, 1912; László *et al.*, 2007; Jiang *et al.*, 2015). *Elsewhere*: Thailand (László *et al.*, 2007; Jiang *et al.*, 2015).

Remark: The subspecies *M. stramineata likiangensis* (Sick, 1941) was described from China, Yunnan (László *et al.*, 2007).

Genus *Nemacerota* Hampson, 1892

Hampson, 1892: *Fauna Br. India (Moths)*, **1**: 185.

Type-species: *Asphalia cinerea* Warren, 1888.

145. *Nemacerota cinerea* (Warren, 1888)

1888. *Asphalia cinerea* Warren, *Proc. Zoo. Soc. Lond.*, **1888**: 300.

2007. *Nemacerota cinerea*: László *et al.*, *Esperiana*, **13**: 124.

Distribution: India: Himachal Pradesh (László *et al.*, 2007). *Elsewhere*: Thandiani, Pakistan (László *et al.*, 2007).

Genus *Neotogaria* Matsumura, 1933

Matsumura, 1933, *Insecta matsum.*, 7 (4): 195.

Type-species: *Neotogaria saitonis* Matsumura, 1931.

146. *Neotogaria anguligera* (Hampson, 1892)

1892. *Polyploca anguligera* Hampson, *Fauna Br. India (Moths)*, **1**: 183.

2007. *Neotogaria anguligera*: László *et al.*, *Esperiana*, **13**: 153.

Distribution: India: Naga hills (Hampson, 1892; László *et al.*, 2007) *Elsewhere*: Thailand (László *et al.*, 2007).

147. *Neotogaria galema* (Swinhoe, 1894)

1894. *Polyploca galema* Swinhoe, *Ann. Mag. Nat. Hist.*, **14**(6): 431.

2007. *Neotogaria galema*: László *et al.*, *Esperiana*, **13**: 152.

Distribution: India: Khasis, Meghalaya, Cherra Punji (Swinhoe, 1894; László *et al.*, 2007). *Elsewhere*: Nepal, Thailand, Vietnam (László *et al.*, 2007).

Remark: László *et al.* (2007) erroneously refer the type locality Cherrapunji as located in Bangladesh. In fact, the historical locality is in the Meghalaya Mts, India.

Genus *Nothoploca* Yoshimoto, 1982

Yoshimoto, 1982; *Tinea*, **11** (4): 125.

Type-species: *Polyploca nigripunctata* Warren, 1915.

148. *Nothoploca nigripunctata nigripunctata* (Warren, 1915)

1915. *Polyploca nigripunctata* Warren, *Novit. Zool.*, **22**(1): 157.

Distribution: India: Sikkim (Warren, 1915; László *et al.*, 2007). *Elsewhere*: Nepal (László *et al.*, 2007).

Remark: Further subspecies are *N. nigripunctata fansipana* László, Ronkay, Ronkay & Witt, 2007 from N. Vietnam, Fansipan and *N. nigripunctata zolotarenski* Dubatolov, 1987 from Russian Far East (László *et al.*, 2007).

Genus *Paragnorima* Warren, 1912

Warren, 1912; In Seitz, *Gross Schmett. Erde*, 2: 329.

Type-species: *Gaurena fuscescens* Hampson, 1892.

149. *Paragnorima fuscescens* (Hampson, 1892)

1892. *Gaurena fuscescens* Hampson, *Fauna Br. India (Moths)*, 1: 182.

1900. *Palimpsestis brunnea* Leech, *Trans., Ento. Soc. Lond.*, 1900: 18.

1921. *Haplothyatira unipunctata* Houlbert, In Oberthür, *Études de Lépidoptérologie Comparée*, 18(2): 115.

1943. *Haplothyatira unipunctata dubitatrix* Bryk, *Ark. Zool.*, 34 A: 13.

2007. *Paragnorima fuscescens*: László et al., *Esperiana* 13: 247.

Distribution: India: Naga Hills (Hampson, 1892; László et al., 2007); Sikkim (Lachen, Lachung) (Houlbert, 1921; László et al., 2007; Jiang et al., 2015). *Elsewhere*: China (Tibet, Yunnan, Sichuan), Nepal, Myanmar, Vietnam (László et al., 2007), Thailand (Jiang et al., 2015).

Genus *Parapsestis* Warren, 1912

Warren, 1912; In Seitz, *Gross Schmett. Erde*, 2: 329.

Type-species: *Cymatophora argenteopicta* Oberthür, 1879.

150. *Parapsestis lichenea lichenea* (Hampson, 1892)

1892. *Gaurena lichenea* Hampson, *Fauna Br. India (Moths)*, 1: 182.

1921. *Cymatophora candida* Houlbert, In Oberthür, *Études de Lépidoptérologie Comparée*, 18(2): 169.

2007. *Parapsestis lichenealichenea*: László et al., *Esperiana*, 13: 242.

Distribution: India: Sikkim, Nagaland (Hampson, 1892; László et al., 2007). *Elsewhere*: Nepal (László et al., 2007).

Remark: Further subspecies are *P. lichenea splendida* László, Ronkay, Ronkay & Witt, 2007 from N. Vietnam, N. Thailand, China-Yunnan and Myanmar and *P. lichenea tsinlinga* László, Ronkay, Ronkay & Witt, 2007 from East China (László et al., 2007).

Genus *Psidopala* Houlbert, 1921

Houlbert, 1921; *Études de Lépidoptérologie Comparee*, 18 (2): 93.

Type-species: *Thyatira opalescens* Alphéraky, 1897.

151. *Psidopala undulans* (Hampson, 1893)

1893. *Thyatira undulans* Hampson, *J. Bomb. Nat. Hist. Soc.*, 1: 283.

1943. *Psidopala ebba* Bryk, *Arkiv för Zoologi*, 34 A: 7.

2007. *Psidopala undulans*: László et al., *Esperiana*, 13: 178.

Distribution: India: Khasis (Hampson, 1893; László et al., 2007). *Elsewhere*: Myanmar (Bryk, 1943; László et al., 2007), China (Tibet) (Jiang et al., 2015).

152. *Psidopala tenuis* (Hampson, 1896)

1896. *Gaurena tenuis* Hampson, *Fauna Br. India (Moths)*, 4: 462.

1966. *Psidopala tenuis falkneri* Werny, *Untersuchungen über die Systematik* (*Lepidoptera, Thyatiridae*): 263.

2007. *Psidopala tenuis*: László et al., *Esperiana*, 13: 186.

Distribution: India: Yatong, Sikkim (Hampson, 1896; László et al., 2007). *Elsewhere*: Nepal (László et al., 2007); China (Tibet) (Jiang et al., 2015).

Genus *Spica* Swinhoe, 1889

Swinhoe, 1889; *Proc. Zool. Soc. Lond.*, 1889: 424.

Type-species: *Spica luteola* Swinhoe, 1889.

153. *Spica luteola* Swinhoe, 1889

1889. *Spica luteola* Swinhoe, *Proc. Zool. Soc. Lond.*, 1889: 424.

Distribution: India: Sikkim (Jiang et al., 2015), Uttarakhand, Joshimath (Smetacek, 2002). *Elsewhere*: Nepal, China, Tibet (Jiang et al., 2015).

Genus *Stenopsestis* Yoshimoto, 1984

Yoshimoto, 1984; *Tyô to Ga*, 34 (1): 27.

Type-species: *Palimpsestis alternata* Moore, 1881.

154. *Stenopsestis alternata* (Moore, 1881)

1881. *Palimpsestis alternata* Moore, *Proc. Zoo. Soc. Lond.*, 1881: 331.

1993. *Stenopsestis alternata bryki* Yoshimoto, 123.

2007. *Stenopsestis alternata*: László et al., *Esperiana*, 13: 244.

Distribution: India: Sikkim, Darjeeling (Moore, 1881; László et al., 2007). *Elsewhere*: China (Sichuan), Nepal, Myanmar, Thailand (László et al., 2007)

Genus *Takapsestis* Matsumura, 1933

Matsumura, 1933; *Insecta Matsumurana* 7: 200.

Type-species: *Takapsestis wilemaniella* Matsumura, 1933.

155. *Takapsestis curvicosta* (Warren, 1915)

1915. *Polyploca curvicosta* Warren, *Novit. zool.*, 22(1): 156.

1990. *Takapsestis curvicosta*: Yoshimoto, *Nota Lepid.*, 13(4): 241.

Distribution: India: Assam, Khasi Hills (Warren, 1915; Yoshimoto, 1990). *Elsewhere*: Nepal (Yoshimoto, 1990); Thailand (László et al., 2007).

156. *Takapsestis bifasciata* (Hampson, 1895)

1895. *Polyploca bifasciata* Hampson, *Trans. Ent. Soc.*, 1895: 281.

1992. *Takapsestis nepalensis* Yoshimoto, *Tinea*, 13(2): 48.

1994. *Takapsestis harutai* Yoshimoto, *Tinea*, 14(1): 92.

1990. *Takapsestis bifasciata*: Yoshimoto, *Nota Lepid.*, 13(4): 236.

Distribution: India: Sikkim (Hampson, 1895; Yoshimoto, 1990; László *et al.*, 2007). *Elsewhere:* Nepal (László *et al.*, 2001; László *et al.*, 2007).

Remark: Year of description of the *Type-species* of the genus was erroneously stated as 1896 by Yoshimoto (1990). Further, *T. nepalensis* and *T. harutai* synonymized by László *et al.* (2001).

157. *Takapsestis orbicularis* (Moore, 1888)

1888. *Polyploca orbicularis* Moore, *Proc. Zoo. Soc. Lond.*, **1888**: 407.

1915. *Polyploca griseata* Warren, *Novit. Zool.*, **22**: 157.

1990. *Takapsestisorbicularis*: Yoshimoto, *Nota Lepid.*, **13**(4): 236.

Distribution: India: Dharmshala, Kangra, Sikkim (Moore, 1888; Hampson, 1892; Warren, 1915), Western Himalaya to Sikkim (Yoshimoto, 1990; László *et al.* 2001). *Elsewhere:* Nepal, Pakistan (László *et al.*, 2001; László *et al.*, 2007).

Remark: *Polyploca griseata* was synonymized by László *et al.* (2001).

Genus *Tethea* Ochseneimer, 1816

Ochseneimer, 1816; *Schmett. Eur.*, **4**: 64.

Type-species: *Noctua* or Denis & Schiffermüller, 1775.

158. *Tethea albicosta* (Moore, 1867)

1867. *Thyatira albicosta* Moore, *Proc. Zool. Soc. Lond.*, 1867: 45.

1966. *Tethea albicosta birmanica* Werny, *Untersuchungen über die Systematik (Lepidoptera, Thyatiridae)*: 394.

2007. *Tethea albicosta*: László *et al.*, *Esperiana*, **13**: 117.

Distribution: India: Shimla, Sikkim (Hampson, 1892; László *et al.*, 2007), Khasi Hills (Moore, 1867; László *et al.*, 2007). *Elsewhere:* China, Nepal, Myanmar, Vietnam (László *et al.*, 2007).

159. *Tethea consimilis commifera* (Warren, 1912)

1912. *Saronaga commifera* Warren, In Seitz, *Gross-Schmett. Erde*, **2**: 322.

1966. *Tethea consimilis szechwanensis* Werny, *Untersuchungen über die Systematik (Lepidoptera, Thyatiridae)*: 407.

2007. *Tethea consimilis commifera*: László *et al.*, *Esperiana*, **13**: 119.

Distribution: India: India (Sabathu) (Warren, 1912; László *et al.*, 2007); Indian Himalayas (Kim *et al.*, 2006), Himachal Pradesh (László *et al.*, 2007). *Elsewhere:* China (Yunnan, Sichuan, Tibet), Nepal (László *et al.*, 2007).

Remarks: The species is widely distributed and represented by five subspecies: the nominate one occurs

in the Pacific region (Japan, Russian Far East, Korea), *T. consimilis c-album* (Matsumura, 1931) is endemic to Taiwan, *T. consimilis commifera* (Warren, 1912) is distributed from Himachal Pradesh through Nepal and Sikkim to Yunnan, Sichuan and the Wuyshan areas in China, *T. consimilis aurisigna* (Bryk, 1943) occurs in the northern regions of Indochina and Malaysia, finally *T. consimilis congener* (Roepke, 1945) is supposedly endemic to Sumatra (László *et al.*, 2007).

160. *Tethea oberthuri oberthuri* (Houlbert, 1921)

1921. *Saronaga oberthuri* Houlbert, *Oberthür Etudes Lep. Comp.*, **18**: 194.

1941. *Tethea oberthuri monticola* Bryk, *Ark. Zool.*, **349**(11): 3.

1966. *Tethea oberthuri occidentalis* Werny, *Untersuchungen über die Systematik (Lepidoptera, Thyatiridae)*: 386.

1966. *Tethea oberthuri chekiangensis* Werny, *ibid.*: 387.

1966. *Tethea oberthuri fukiensis* Werny, *ibid.*: 388.

2007. *Tethea oberthuri oberthuri*: László *et al.*, *Esperiana*, **13**: 115.

Distribution: India: N.E. Himalaya (Holloway, 1998); Sikkim, Lachoong and Darjeeling (Werny, 1966; László *et al.*, 2007; Zhuang *et al.*, 2017). *Elsewhere:* Thailand, China, Nepal, Myanmar, Vietnam, Malaysia. (László *et al.*, 2007; Zhuang *et al.*, 2017).

Remarks: Further subspecies are *T. oberthuri taiwana* (Matsumura, 1931) from Taiwan and *T. oberthuri baluensis* Werny, 1966 from Borneo (László *et al.*, 2007).

Genus *Thyatira* Ochseneimer, 1816

Ochseneimer, 1816; *Schmett. Eur.* **4**: 77.

Type-species: *Phalaena batis* Linnaeus, 1758.

161. *Thyatira batis rubrescens* (Werny, 1966)

1966. *Thyatira rubrescens* Werny, *Untersuchungen über die Systematik (Lepidoptera, Thyatiridae)*: 36.

1966. *Thyatira rubrescens nepalensis* Werny, *ibid.*, 38.

1966. *Thyatira rubrescens obscura* Werny, *ibid.*, 39.

1966. *Thyatira rubrescens assamensis* Werny, *ibid.*, 41.

1966. *Thyatira rubrescens szechwana* Werny, *ibid.*, 42.

1966. *Thyatira rubrescens orientalis* Werny, *ibid.*, 42.

1966. *Thyatira rubrescens kwangtungensis* Werny, *ibid.*, 43.

1966. *Thyatira rubrescens tienmushana* Werny, *ibid.*, 44.

1966. *Thyatira rubrescens vietnamensis* Werny, *ibid.*, 48.

1992. *Thyatira batis rubrescens*: Yoshimoto, *Tinea*, **13**(2): 47.

Distribution: India: Sikkim, Assam, Cherrapunji (Werny, 1966; Yoshimoto, 1992; László *et al.*, 2007; Jiang *et al.*, 2015; Zhuang *et al.*, 2017), Kumaon, Nainital (László

et al., 2007). *Elsewhere*: China, Nepal, Vietnam, Thailand (Werny, 1966; Yoshimoto, 1992; László et al., 2007; Jiang et al., 2015; Zhuang et al., 2017).

Remark: *Thyatira batis* is a widely distributed taxon in Eurasia with the following other subspecies: *T. batis batis* (Linnaeus, 1758)(Russia, N-NE China, Europe), *T. batis formosicola* Matsumura, 1933 (Taiwan) and *T. batis pallida* Rothschild, 1920 (Sumatra) (László et al., 2007).

162. *Thyatira cognata* Warren, 1888

1888. *Thyatira cognata* Warren, *Proc. Zoo. Soc. Lond.*, 1888: 299.

1888. *Thyatira cognata* Moore, *Proc. Zoo. Soc. Lond.*, 1888: 406. (László et al., 2007; *Esperiana*, 13: 46)

Distribution: India: Himachal Pradesh, Kangra (Warren, 1888; Moore, 1888; Werny, 1966), Kashmir (Srinagar), Himachal Pradesh (Lahoul) (László et al., 2007). *Elsewhere*: Pakistan (László et al., 2007).

Remark: Due to page priority, the species' first author is Warren (László et al., 2007).

Genus *Toxoides* Hampson, 1892

Hampson, 1892; *Fauna Br. India (Moths)*, 1: 185.

Type-species: *Toxoides longipennis* Hampson, 1892.

163. *Toxoides undulatus* (Moore, 1867)

1867. *Osica undulatus* Moore, *Proc. Zool. Soc. Lond.*, 1867: 45.

1892. *Toxoides longipennis* Hampson, *Fauna Br. India (Moths)*, 1: 186.

1943. *Toxoides emphloioides* Bryk, *Ark. Zool.*, 34 A (11): 15.

1992. *Toxoides undulatus*: Yoshimoto, *Tinea*, 13(2): 49.

Distribution: India: Khasi Hills, Sikkim, Darjeeling, Tiger Hill (Moore, 1867; Hampson, 1892, Yoshimoto, 1992). *Elsewhere*: Myanmar (Bryk, 1943), China (Yunnan), Nepal (Yoshimoto, 1992). Vietnam, Thailand (László et al., 2007).

References

- Bryk, F. 1943. Entomological results from the Swedish expedition 1934 to Burma and British India. Lepidoptera: Drepanidae. *Ark. Zool.*, 34 A (13): 1-30.
- Buchsbau, U. 2010. *Nordstromia guenterriedeli* sp. n. from Sumatra (Indonesia) (Lepidoptera, Drepanidae). *Entomofauna*. 31 (7): 57-68.
- Butler, A. G. 1886. Illustration of typical specimens of Lepidoptera heterocera in the collection of the British Museum. 6: 1-89.
- Butler, A. G. 1889. Illustration of typical specimens of Lepidoptera heterocera in the collection of the British Museum. 7: 1-124.
- Chandra, K. and Sambath, S. 2013. Moth diversity of Tawang District, Arunachal Pradesh, India. *Journal of Threatened Taxa*. 5 (1): 3565-3570. <https://doi.org/10.11609/JOTT.02718.966>.
- Cotes, E. C. and Swinhoe, C. C. (1887-1889). *A Catalogue of Moths of India. Part IVI: Sphinges, Bombyces, Noctues, Pseudo- Deltoids and Deltoids, Gephyrites, Pyrales, Crambites, Tortrices and Addenda*, Calcutta; p. 1-812.
- Dey, P. and Sondhi, S. 2019. First records of *Agnidra vinacea* (Moore, 1879) (Lepidoptera: Drepanidae: Drepaninae) from the western Himalaya, extending its known range westwards. *Journal of Threatened Taxa*. 11 (5): 13622-13624. <https://doi.org/10.11609/>

Genus *Wernya* Yoshimoto, 1987

Yoshimoto, 1987; *Tyô to Ga*, 38(1): 39.

Type-species: *Palimpsestis lineofracta* Houlbert, 1921.

164. *Wernya solena* (Swinhoe, 1894)

1894. *Gaurena solena* Swinhoe, *Ann. Mag. Nat. Hist.*, 14(6): 431.

1987. *Wernya solena*: Yoshimoto, *Tyo to Ga*, 38(1): 47.

Distribution: India: NE India, Assam, Khasis (Yoshimoto, 1987; László et al., 2007). *Elsewhere*: not known.

Discussion

The two families, Drepanidae and Geometridae were considered united in the old concept of superfamily Geometroidea. Presently, both are considered as well distinct groups representing separate superfamilies: Drepanoidea and Geometroidea. Family Geometridae from India was compiled by Kirti et al. (2019) and is known by 2041 species but the data was lacking about the current status of Drepanidae. This updated checklist of 164 species (including subspecies) under 55 genera of family Drepanidae presented here will fill that gap. The detailed information about distribution of the species along with complete synonymy will act as baseline for lepidopterist to initiate work on family Drepanidae from India.

Acknowledgements

Authors are thankful to Dr. Kailash Chandra, Director, Zoological Survey of India (ZSI) for providing his guidance and encouragement to compile this work. Some short notes provided by Dr J.D. Holloway (NHM, London) and some important literature on Drepanoidea provided by Mrs. Alka Vaidya (Mumbai) and Dr Joel Minet (France) to first author is also duly acknowledged.

- jott.4593.11.5.13622-13624.
- Freina, de. 1994. Beitrag zur systematischen Erfassung der Bombyces- und Sphinges Fauna Kleinasiens. Weitere Kenntnisse über Artenspektrum, Systematik und Verbreitung von Cossidae, Psychidae, Cochlidiidae, Syntomidae, Saturniidae, Brahmaeidae, Drepanidae, Axiidae, Hepialidae, Dilobidae und Nolidae (Insecta, Lepidoptera) *Atalanta*. **25** (1/2): 317-349.
- Gaede, M. 1931. Drepanidae. In: Strand, E.H. (Ed.), *Lepidopterorum Catalogus*, Berlin; **49**:1-59.
- Hampson, G. F. 1892. *The Fauna of British India, Moths*. 1. Taylor and Francis, London, British; p. xxiii+527.
- Hampson, G. F. 1893. Illustrations of Typical Specimens of Lepidoptera Heterocera in the Collection of the British Museum, Part IX. The Macrolepidoptera Heterocera of Ceylon. Taylor and Francis, London; p. 182.
- Hampson, G. F. 1895. Descriptions of new Heterocera from India. *Transactions of the Entomological Society of London*: 277-315. <https://doi.org/10.1111/j.1365-2311.1895.tb01672.x>
- Hampson, G. F. 1896. *The Fauna of British India, Moths*. 4. Taylor and Francis, London, British; p. xxviii + 594.
- Hampson, G. F. 1912. The moths of India. Supplementary paper to the volumes in 'The Fauna of British India'. Series 4, Part 6. *J. Bombay Nat. Hist. Soc.* **21**: 1222-1272.
- Hampson, G. F. 1914. Descriptions of new genera and species of Drepanidae and Thyrididae. *Ann. Mag. Nat. Hist.* **14** (8): 103-117. <https://doi.org/10.1080/00222931408693551>.
- Holloway, J. D. 1998. The Moths of Borneo: Part 8; Family Castniidae, Callidulidae, Drepanidae, Uraniidae. *Malayan Nature Journal*. **52**: 1-155.
- Holloway, J. D. 2011. The Moths of Borneo (part 2): families Phaudidae, Himantopteridae and Zygaenidae; revised and annotated checklist. *Malayan Nature Journal*. **63**: 1-548.
- Holloway, J. D., Kirby, G. and Peggie, D. 2001. *Fauna Malesiana Handbooks: The Families of Malesian Moths and Butterflies*. Leiden, The Netherlands, Brill; p. 455.
- Imms, A. 1934. *A General Textbook of Entomology*. London, Methuen & Co; p. xii+727.
- Inoue, H. 1988. Taxonomic notes on the Drepanidae from Taiwan, with descriptions of two new species and six new subspecies (Lepidoptera). *Tinea*. **12**: 125-134.
- Jiang, N., Yang, C., Xue, D. and Hang, H. 2015. An updated checklist of Thyatirinae (Lepidoptera, Drepanidae) from China, with descriptions of one new species. *Zootaxa*. **3941**(1): 1-48. <https://doi.org/10.11646/zootaxa.3941.1.1>. PMID: 25947492.
- Jiang, N., Liu, S., Xue, D. and Han, H. 2016. A review of Cyclidiinae from China (Lepidoptera, Drepanidae). *ZooKeys*. **553**: 119-148. <https://doi.org/10.3897/zookeys.553.6153>. PMID: 26877675, PMCID: PMC4740984.
- Kim, M.Y., Lee H.K., Ronkay L. and Park, K.T. 2006. A Review of the Korean Thyatiridae (Lepidoptera), including the Mt. Changbaisan. *J. Asia-Pacific Entomol.*, **9** (3): 203-221. [https://doi.org/10.1016/S1226-8615\(08\)60293-9](https://doi.org/10.1016/S1226-8615(08)60293-9).
- Kirti, J. S., Chandra, K., Saxena, A. and Singh, N. 2019. *Geometrid moths of India*. Nature Books India, New Delhi; p. 1-296.
- Kristensen, N. P., Scoble, M. J. and Karsholt, O. 2007. Lepidoptera phylogeny and systematics: The State of inventorying moth and butterfly diversity. *Zootaxa*. **1668**: 699-747. <https://doi.org/10.11646/zootaxa.1668.1.30>.
- László, G. M. and Ronkay, G. 2000. New *Epipsestis* Matsumura, 1921 species (Lepidoptera, Thyatiridae) from Vietnam and from Nepal. *Acta Zoologica Academiae Scientiarum Hungaricae*. **46** (4): 337-349.
- László, G. M., Ronkay, G. and Ronkay, L. 2001. Taxonomic studies on the Eurasian Thyatiridae. Revision of Wernya Yoshimoto, 1987 generic complex and the genus *Takapsestis* Matsumura, 1933 (Lepidoptera). *Acta Zoologica Academiae Scientiarum Hungarica*. **47** (1): 27-85.
- László, G. M., Ronkay, G., Ronkay, L. and Witt, T. 2007. The Thyatiridae of Eurasia: including the Sundaland and New Guinea (Lepidoptera). *Esperiana*. **13**: 2-683.
- Matsumura, S. 1921. *Thousand Insects of Japan Additamenta*. Tokyo. **4**: 772-1012,
- Matsumura, S. 1927. New species and subspecies of moths from the Japanese Empire. *Coll. Agric. Hokkaido Imp. Univ.* **19**: 1-91.
- Matsumura, S. 1931. 6,000 Illustrated Insects of Japan-Empire. Tokyo. **1497** -f-: 191.
- McDunnough, J. 1938. Check List of the Lepidoptera of Canada and the United States of America. Part I. Macrolepidoptera. *Memoirs of Southern California Academy of Sciences*, 1: 1-275.
- Minet, J. 1983. Étude morphologique et phylogénétique des organes tympaniques des Pyra-loidea. I. généralités et homologies. (Lep. Glossata). *Annales de la Société entomologique de France (N.S.)*. **19**: 175-207 [In French].
- Minet, J. 1991. Tentative reconstruction of the ditrysian phylogeny (Lepidoptera: Glossata). *Entomologica Scandinavica*. **22**: 69-95. <https://doi.org/10.1163/187631291X00327>.
- Minet, J. 2002. The Epicopeiidae: Phylogeny and a redefinition, with the description of new taxa (Lepidoptera: Drepanoidea). *Annales de la Societe Entomologique de France*. **38**: 463-487. <https://doi.org/10.1080/00379271.2002.10697355>.
- Minet, J. and Scoble, M. J. 1999. The drepanoid/geometroid Assemblage. In: Kristensen NP (Ed.) *Lepidoptera, Moths and Butterflies*. Vol. I: Evolution, Systematics, and Biogeography. Handbook of Zoology. Vol. IV, Arthropoda: Insecta, Part 35. Walter de Gruyter, Berlin and New York; p. 301-320.
- Moore, F. 1866. On the Lepidopterous Insects of Bengal. *Proceedings Zoological Society London*. **1865**: 755-823. <https://doi.org/10.1111/j.1469-7998.1865.tb02432.x>.

- Moore, F. T. and Moore, F., 1858. *A Catalogue of the Lepidopterous Insects in the Museum of the Honourable East India Company*. **2**: 279-438.
- Moore, F. 1879. Descriptions of new Indian lepidopterous insects from the collection of the late Mr. W. S. Atkinson. **1**: 1-88.
- Moore, F. 1887. Descriptions of new Indian lepidopterous insects from the collection of the late Mr. W. S. Atkinson. **3**: 199-299.
- Moore, F. 1888. Descriptions of new Genera and Species of Lepidoptera Heterocera collected by Rev. J. H. Hocking, chiefly in the Kangra District, N.W. Himalaya. *Proceedings Zoological Society London*; **1888**: 390-412. <https://doi.org/10.1111/j.1469-7998.1888.tb06720.x>.
- Mutanen, M., Wahlberg, N. and Lauri, K. 2010. Comprehensive gene and taxon coverage elucidates radiation patterns in moths and butterflies. *Proceedings Royal Society B*. **277**: 2839-2848. <https://doi.org/10.1098/rspb.2010.0392>. PMID: 20444718, PMCID: PMC2981981.
- Nakamura, M. 1981. Key to the classification of the Japanese lepidopterous pupae. *Tyô to Ga*. **32**: 1-12.
- Nieukerken, E.J., Kaila, L., et al. Order Lepidoptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa*. **3148**: 212-221. <https://doi.org/10.11646/zootaxa.3148.1.41>.
- Regier, J.C., Zwick, A., Cummings, M.P., et al. 2009. Toward reconstructing the evolution of advanced moths and butterflies (Lepidoptera: Ditrysia): An initial molecular study. *BMC Evolutionary Biology*. **9**: 280. doi:10.1186/1471-2148-9-280. <https://doi.org/10.1186/1471-2148-9-280>. PMID: 19954545, PMCID: PMC2796670.
- Sanyal, A.K., Mallick, K., Khan, S., Bandyopadhyay, U., Mazumder, A., Bhattacharyya, K., Pathania, P.C., Raha, A. and Chandra, K. 2018. Insecta: Lepidoptera (Moths). In: *Faunal Diversity of Indian Himalaya*. p. 651-726. (Published by the Director, Zool. Surv. India, Kolkata).
- Scoble, M. J. 1992. *The Lepidoptera: Form, Function, and Diversity*. Oxford University Press; p. 1-404.
- Smetacek, P. 2002. Notes on new records of hooktip moths, Lepidoptera: Drepanidae, from the Kumaon and Garhwal Himalaya. *J. Bombay Nat. Hist. Soc.*, **99** (3): 446-454.
- Sondhi, Y. and Sondhi S. 2016. A partial checklist of moths (Lepidoptera) of Dehradun, Mussoorie and Devalsari in Garhwal, Uttarakhand, India. *Journal of Threatened Taxa*. **8** (5): 8756-8776. <https://doi.org/10.11609/jott.2814.8.5.8756-8776>.
- Song, W. H., Xue, D. Y. and Han, H. X. 2011. A taxonomic revision of *Tridrepana* Swinhoe, 1895 in China, with descriptions of three new species (Lepidoptera, Drepanidae). *Zootaxa*. 3021: 39-62.
- Song, W. H., Xue, D. Y. and Han, H. X. 2012 Revision of Chinese Oretinae (Lepidoptera, Drepanidae). *Zootaxa*. **3445**: 1-36. <https://doi.org/10.11646/zootaxa.3445.1.1>.
- Strand, E. 1911. Eucherinae. In: Seitz A, ed. *Die Gross-Schmetterlinge der Erde*, 2; p. 195-206.
- Swinhoe, 1894. New species of Eastern Lepidoptera. *The Annals and Magazine of Natural History*. (6) **14** (84): 429-443. <https://doi.org/10.1080/00222939408677830>.
- Swinhoe, C. 1899. New species of Oriental Lepidoptera. *The Annals and Magazine of Natural History*. **7**: 102-116. <https://doi.org/10.1080/00222939908678084>.
- Swinhoe, C. 1902. New and little known species of Drepanulidae, Epiplemidae, Microniidae and Geometridae in the national collection. *Transactions of the Royal Entomological Society of London*. **3**: 584-677.
- Swinhoe, C. 1905. Notes on Eastern and Australian Heterocera, with descriptions of one new genus and thirteen new species. *Annals and Magazine of Natural History*. **16** (7): 142-155. <https://doi.org/10.1080/03745480509443661>.
- Walker, F. 1860. List of the specimens of Lepidopterous Insects in the collection of the British Museum. Edward Newman, London. **20**: 1-276.
- Walker, F. 1862. List of the specimens of Lepidopterous Insects in the collection of the British Museum. Edward Newman, London. **24**: 1021-1280.
- Walker, F. 1862a. List of the specimens of Lepidopterous Insects in the collection of the British Museum. Edward Newman, London. **25**: 1281-1477.
- Walker, F. 1866. List of the specimens of Lepidopterous Insects in the collection of the British Museum. Edward Newman, London. **35**: 1535-2040.
- Warren, W. 1896. New species of Drepanulidae, Thyrididae, Uraniidae, Epiplemidae and Geometridae in the Tring Museum. *Novitates Zoologicae*. **3**: 335-419. <https://doi.org/10.5962/bhl.part.5427>.
- Warren, W. 1897. New genera and species of moths from the Old World regions in the Tring Museum. *Novitates Zoologicae*. **4**: 12-130. <https://doi.org/10.5962/bhl.part.21182>.
- Warren, W. 1901. Drepanulidae, Uraniidae, and Geometridae from the Palaearctic and Indo- Australian Regions. *Novitates Zoologicae*. **8**: 190-201.
- Warren, W. 1902. Drepanulidae, Thyrididae, Uraniidae, Epiplemidae and Geometridae from the Oriental region. *Novitates Zoologicae*. **9**: 340-372.
- Warren, W. 1914. New species of Drepanulidae, Noctuidae and Geometridae in the Tring Museum. *Novitates Zoologicae*. **21**: 401-425. <https://doi.org/10.5962/bhl.part.16333>.

- Warren, W. 1922. Family: Drepanidae. In: Seitz A (Ed.) *The Macrolepidoptera of the World*. Vol. 10: Bombyces and Sphinges of the Indo-Australian Region. Alfred Kernen, Stuttgart; p. 443-472.
- Warren, W. 1923. Family: Drepanidae. In: Seitz A (Ed.), *The Macrolepidoptera of the World*. Vol. 10: Bombyces and Sphinges of the Indo-Australian Region. Alfred Kernen, Stuttgart; p. 473-490.
- Watson, A. 1957. A revision of the genus *Tridrepana* Swinhoe (Lepidoptera: Drepanidae). *Bull. Br. Mus. Nat. Hist. (Ent.)*. **4**: 409-500.
- Watson, A. 1957a. A Revision of the genus *Deroca* Walker (Lepidoptera, Drepanidae). *Ann. Mag. Nat. Hist.* **10** (12): 129-148. <https://doi.org/10.1080/00222935708655939>.
- Watson, A. 1959. A Revision of the genus *Auzata* Walker (Lepidoptera, Drepanidae). *Bonn. zool. Beitr.* **9**: 232-257.
- Watson, A. 1961. A taxonomic study of some Indo-Australian Drepanidae (Lepidoptera) *Bull. Br. Mus. nat. Hist. (Ent.)*. **10**: 317-347. <https://doi.org/10.5962/bhl.part.16263>.
- Watson, A. 1965. A Revision of the Ethiopian Drepanidae (Lepidoptera). *Bull. Br. Mus. Nat. Hist. (Ent.)*. Suppl. **3**: 1-178 pp.
- Watson, A. 1967. A survey of the extra-Ethiopian Oretinae (Lepidoptera: Drepanidae). *Bull. Br. Mus. Nat. Hist. (Ent.)*. **19**: 151-221. <https://doi.org/10.5962/bhl.part.28301>.
- Watson, A. 1968. The taxonomy of the Drepaninae represented in China, with an account of their world distribution (Lepidoptera: Drepanidae). *Bull. Br. Mus. Nat. Hist. (Ent.)*. Supplement **12**: 1-151.
- Werny, K. 1966. *Untersuchen über die Systematik der Tribus Thyatirini, Macrothyatirini, Habrosynini und Tetheini (Lepidoptera: Thyatiridae)*. Inaugural-Dissertation, Universität des Saarlandes; p. 1-463.
- Wilkinson, C. 1967. A taxonomic revision of the genus *Teldenia* Moore (Lepidoptera: Drepanidae, Drepaninae). *Trans. R. Ent. Soc. Lond.* **119**: 303-362. <https://doi.org/10.1111/j.1365-2311.1967.tb00502.x>.
- Wilkinson, C. 1968. A taxonomic revision of the genus *Ditrigona* (Lepidoptera: Drepanidae: Drepaninae). *Trans. Zool. Soc. Land.* **31**: 403-517. <https://doi.org/10.1111/j.1096-3642.1968.tb00369.x>.
- Wilkinson, C. 1970. On the taxonomy of *Dipriodontia sericea* Warren (Lepidoptera: Drepanidae). *Proc. R. Ent. Soc. Lond. (B)*. **39**: 89-98. <https://doi.org/10.1111/j.1365-3113.1970.tb00261.x>.
- Wilkinson, C. 1972. *The Drepanidae of Nepal (Lepidoptera)*. Khumbu Himal, Ergebnisse Forschungsunternehmens Nepal Himalaya. **4**: 157-332.
- Wu, C. G., Han, H. X. and Xue, D. Y. 2010. A pilot study on the molecular phylogeny of Drepanoidea (Insecta: Lepidoptera) inferred from the nuclear gene EF-1a and the mitochondrial gene COI. *Bulletin of Entomological Research*. **100**: 207-216. <https://doi.org/10.1017/S0007485309990162>. PMID: 19580687.
- Yazaki, K. 1992. Drepanidae. Moths of Nepal, Part 1, *Tinea*. **13** (Suppl. 2): 1-3.
- Yazaki, K. 1993. Drepanidae. Moths of Nepal, Part 2, *Tinea*. **13** (Suppl. 3): 115-121.
- Yoshimoto, H. 1982. Notes on the genus *Epipsestis*, with descriptions of three new species from Nepal (Lepidoptera, Thyatiridae). *Tyô To Ga*. **32** (3-4), 117-137.
- Yoshimoto, H. 1984. A revision of the genus *Demopsestis* and its three related genera, with description of a new species from Taiwan (Lepidoptera, Thyatiridae). *Tyô To Ga*. **34**(1): 1-20.
- Yoshimoto, H. 1984a. Notes on *Epipsestis* species (Thyatiridae) from Taiwan, with description of a new subspecies. *Japan Heterocerists' Journal*. **128**: 33-34.
- Yoshimoto, H. 1987. Notes on *Mimopsestis* Matsumura, 1921, and its allied new genus, with descriptions of three new species from Southeast Asia (Lepidoptera, Thyatiridae). *Tyô to Ga*. **38**: 39-53.
- Yoshimoto, H. 1990. *Takapsestis fascinâta* sp. n. from China (Lepidoptera, Thyatiridae). *Nota lepid.*, **13**(4): 236-241
- Yoshimoto, H. 1992. Thyatiridae. Moths of Nepal, Part 1, *Tinea*, **13**. (Suppl. 2): 47-49.
- Yoshimoto, H. 1993. Thyatiridae, Moths of Nepal, Part 2, *Tinea*, **13**. (Suppl. 3): 122-123.
- Zhuang, H., Yago, M., Owada, M. and Wang, M. 2017. Taxonomic review of the moth family Thyatiridae (Lepidoptera) from Yunnan province, China. *Zootaxa*. **4306**(4): 451-477. <https://doi.org/10.11646/zootaxa.4306.4.1>.