

Radiation effect of an unsteady MHD free convective flow past an infinite vertical plate with heat, mass transfer and constant suction

¹N. Senapati, ²R.K.Dhal and ^{3*}T.K. Das

¹Department of Mathematics, Ravenshaw University, Cuttack, Orissa-753003, India

²J.N.V. Paralakhemundi, Gajapati, Orissa-761201, India

³Gurukul institute of technology, BBSR, Orissa, India

Email:¹nityananda.senapati@gmail.com,²dhal.rajendra@gmail.com, ^{3*}tusharkant.dash078@gmail.com

*Corresponding author: Dr.T.K. Das, Gurukul Institute of Technology, BBSR, Orissa, India, Email: tusharkant.dash078@gmail.com

Abstract

Free convection MHD flow of a viscous incompressible fluid past an infinite vertical plate with heat and mass in the presence of radiation has been studied. The dimensionless governing equations are solved using Laplace transfer technique. The plate temperature and mass concentrations are raised harmonically. The result are obtained for velocity, temperature, concentration, Nusselt number, Sherwood number and skin friction. The effect of various material parameters are discussed on flow variable and presented by graphs.

Keywords: Contourlet transform; Maximum likelihood detector; Multiplicative image watermarking.

Introduction

Free convection flows past different types of vertical bodies are studied because of their wide applications and hence, it has attracted the attention of numerous investigators and scientists. Literature on unsteady MHD convection heat transfer with or without Hall currents are very extensive due to its technical importance in the scientific community (Elbashbeshy, 1988; Rohsenow, 1998). The radiation effects on MHD flow and heat transfer problems have become more important industrially. At high operating temperature, radiation effect can be quite significant (Sahoo, 2003). Many processes in engineering areas occur at high temperatures and knowledge of radiation heat transfer becomes very important for the design of the pertinent equipment. Nuclear power plants, gas turbines and the various propulsion devices for aircraft, missiles, satellites, and space vehicles are examples of such engineering areas.

Heat and mass transfer of a moving vertical plate with suction was studied by Erickson et al, (1966) and (Gupta & Gupta, 1977) with different conditions. Convective heat transfer in an electrically conducting fluid at stretching surface

was studied by (Vajravelu & Hadjinicolaou, 1977). Unsteady free convection flow past a vertical porous plate was investigated by (Helmy, 1988). Acharya *et al.* (2000) have studied free convection and mass transfer flow through a porous medium bounded by vertical infinite surface with constant suction and heat flux (Chen & Char, 1998). But in those studies, they considered the flow to be steady (Danberg & Fansber 1976). Coming back to unsteady case, (Young, 2000), investigated unsteady MHD convective heat transfer past a semi-infinite vertical porous moving plate with variable suction. (Chamka, 2004), considered the case taking mass transfer with heat absorption (Sakiadis, 1961). Muthukumaraswamy & Kumar, (2004) investigated heat and mass transfer effect on moving vertical plate in the presence of Thermal Radiation (Carnahan, 1969; Takhar *et al.*, 1996; Singh *et al.*, 2003). In this paper, we have discussed free convection MHD flow of a viscous incompressible fluid past an infinite vertical plate with heat and mass in the presence of radiation.

Formulation of the Problem

Consider an unsteady two dimensional free convective flow of an electrically conducting,

radiating, viscous and incompressible fluid past an infinite vertical porous plate with constant suction. X_1 -axis is taken along the vertical plate in upward direction and the Y_1 -axis is taken normal to the plate in the direction of the applied uniform magnetic field of strength H_0 . The magnetic permeability is constant throughout the field. There exists a free convection current in the vicinity of the plate. It is assumed that a fluid has constant properties and the variation of density with temperature and mass concentration is considered only in the body force term. All the variable in this flow are the function of y_1 and time t_1 only as the

plate is infinite length. Initially, the temperature and mass concentration at the plate are respectively T_p and C_p and the temperature and mass concentration of fluid are respectively T_∞ and C_∞ . At time $t_1 > 0$, it is assumed that the temperature and mass concentration at the plate rise to $T_1 = T_p + \epsilon(T_p - T_\infty)e^{i\omega_1 t_1}$ and $C_1 = C_p + \epsilon(C_p - C_\infty)e^{i\omega_1 t_1}$. The fully developed flow of a radiating gas by usual Boussinesq's approximation the unsteady flow governed by following equation.

$$\frac{\partial v_1}{\partial y_1} = 0 \tag{1}$$

$$\frac{\partial u_1}{\partial t_1} + v_1 \frac{\partial u_1}{\partial y_1} = g\beta(T_1 - T_\infty) + g\beta_c(C_1 - C_\infty) + \nu \frac{\partial^2 u_1}{\partial y_1^2} - \frac{\sigma B_0^2 u_1}{\rho} \tag{2}$$

$$\frac{\partial T_1}{\partial t_1} + v_1 \frac{\partial T_1}{\partial y_1} = k \frac{\partial^2 T_1}{\partial y_1^2} - \frac{\partial q_r}{\partial y_1} \tag{3}$$

$$\frac{\partial C_1}{\partial t_1} + v_1 \frac{\partial C_1}{\partial y_1} = D' \frac{\partial^2 C_1}{\partial y_1^2} \tag{4}$$

The initial and boundary conditions of the problem are

$$t_1 \leq 0 : u_1 = 0 \quad T_1 = T_\infty \quad C_1 = C_\infty \quad \text{for all } y_1$$

$$t_1 > 0 :$$

$$u_1 = V_0, \quad T_1 = T_p + \epsilon(T_p - T_\infty)e^{i\omega_1 t_1},$$

$$C_1 = C_p + \epsilon(C_p - C_\infty)e^{i\omega_1 t_1} \quad \text{at } y_1 = 0 \tag{5}$$

$$u_1 \rightarrow 0, T_1 \rightarrow T_\infty, C_1 \rightarrow C_\infty \quad \text{as } y_1 \rightarrow \infty$$

where ρ is the density, g acceleration due to gravity, β is the co-efficient of thermal expansion, k is the thermal conductivity, q_r the radiation flux, ν the kinematic viscosity, σ is electrical conductivity,

$B_0 (= H_0 \mu e)$ is the electromagnetic induction. β_c is the co-efficient of expansion of mass and D is the diffusion constant.

Since the plate is assumed to be porous and through it suction with uniform velocity occurs, Equation (1) integrates to $v_1 = -V_0$ is the constant suction velocity

In an optically thick limit, the fluid does not absorb its own emitted radiation, that is, there is no self absorption, but it does absorb radiation emitted by the boundaries. It has been shown by Cogly *et al.* [2]

$$\frac{\partial q_r}{\partial y} = 4(T_1 - T_\infty) \int_0^\infty k_{\lambda w} \left(\frac{de_{b\lambda}}{dT} \right) d\lambda = 4I(T_1 - T_\infty) \quad (6)$$

Where $K_{\lambda w}$ is the absorption co-efficient, $e_{b\lambda}$ is the Plank's function and subscription w refers the value of the wall (plate).

On introducing the following non-dimensional quantities

$$\left. \begin{aligned} t &= \frac{t_1 V_0^2}{4\nu}, y = \frac{y_1 V_0}{\nu}, u = \frac{u_1}{V_0}, Pr = \frac{\nu}{k}, Sc = \frac{\nu}{D} \\ T &= \frac{T_1 - T_\infty}{T_p - T_\infty}, C = \frac{c_1 - c_\infty}{c_p - c_\infty}, \omega = \frac{4\nu\omega_1}{V_0^2}, N = \frac{16I\nu}{V_0^2} \\ M &= \frac{\nu B_0^2 \sigma}{\rho V_0^3}, Gr = \frac{\nu g \beta (T_p - T_\infty)}{V_0^3}, Gm = \frac{\nu g \beta_c (c_p - c_\infty)}{V_0^3} \end{aligned} \right\} \quad (7)$$

where Gr is Grashof number, Gm modified Grashof number, M is magnetic number, Pr is Prandtl number, Sc is Schmidt number, T is non-dimensional temperature, C is non-dimensional mass concentration, ω is oscillating frequency and N is the radiation parameter.

With the help of equation (7) the equations (2) to (5) reduce to

$$\frac{\partial u}{\partial t} - 4 \frac{\partial u}{\partial y} = 4 \frac{\partial^2 u}{\partial y^2} + 4GrT + 4GmC - 4Mu \quad (8)$$

$$\frac{\partial T}{\partial t} - 4 \frac{\partial T}{\partial y} = \frac{4}{Pr} \frac{\partial^2 T}{\partial y^2} - NT \quad (9)$$

$$\frac{\partial C}{\partial t} - 4 \frac{\partial C}{\partial y} = \frac{4}{Sc} \frac{\partial^2 C}{\partial y^2} \quad (10)$$

With the following boundary conditions

$$u = 0, T = 0, C = 0 \text{ for all } y = 0 \text{ and } t \leq 0$$

$$t > 0 : u = 0, T = 1 + \epsilon e^{it\omega}, C = 1 + \epsilon e^{it\omega} \text{ at } y = 0 \quad (11)$$

$$u = 0, T = 0, C = 0 \quad \text{as } y \rightarrow \infty$$

Method of solution

We solve the governing equations in an exact form with the help of using Laplace transforms of equations (8) to (10) using condition (11), we have

$$4 \frac{d^2 \bar{u}}{dy^2} + 4 \frac{d\bar{u}}{dy} - (s + 4M)\bar{u} = 4Gr\bar{T} + 4Gm\bar{C} \quad (12)$$

$$4 \frac{d^2 \bar{T}}{dy^2} + 4Pr \frac{d\bar{T}}{dy} - (NPr + Pr.s)\bar{T} = 0 \quad (13)$$

$$4 \frac{d^2 \bar{C}}{dy^2} + 4Sc \frac{d\bar{C}}{dy} - Sc.s\bar{C} = 0 \quad (14)$$

where s is the Laplace transform parameter

The boundary conditions reduce to

$$\left. \begin{aligned} \bar{u} = 0, \bar{T} = \frac{1}{s} + \frac{\epsilon}{s - \omega i}, \bar{C} = \frac{1}{s} + \frac{\epsilon}{s - \omega i} \text{ at } \bar{y} = 0 \\ \bar{u} = 0, \bar{T} = 0, \bar{C} = 0 \quad \text{as } \bar{y} \rightarrow \infty \end{aligned} \right\} \quad (15)$$

By solving equations (12) to (14) using (15), we get

$$\bar{u} = - \left\{ 4Gm \left(\frac{1}{s} + \frac{\epsilon}{s - \omega i} \right) \frac{1}{(Sc - 1)s - 4M} + 4Gr \left(\frac{1}{s} + \frac{\epsilon}{s - \omega i} \right) \frac{1}{(Pr - 1)s - (4M - NPr)} \right\} e^{-\frac{(1 + \sqrt{1 + s + 4M})y}{2}}$$

$$+ 4Gm \left(\frac{1}{s} + \frac{\epsilon}{s - \omega i} \right) \frac{1}{(Sc - 1)s - 4M} e^{-\frac{(Sc + \sqrt{Sc^2 + Sc.s})y}{2}} \quad (16)$$

$$+ 4Gr \left(\frac{1}{s} + \frac{\epsilon}{s - \omega i} \right) \frac{1}{(Pr - 1)s - (4M - NPr)} e^{-\frac{(Pr + \sqrt{Pr^2 + NPr + Pr.s})y}{2}} \quad (16)$$

$$\bar{T} = \left(\frac{1}{s} + \frac{\epsilon}{s - \omega i} \right) e^{-(Pr + \sqrt{Pr^2 + NPr + Pr.s})y/2} \quad (17)$$

$$\bar{C} = \left(\frac{1}{s} + \frac{\epsilon}{s - \omega i} \right) e^{-(Sc + \sqrt{Sc^2 + Sc.s})y/2} \quad (18)$$

Again by taking inverse Laplace transform of equations (16) – (18), we get

$$\begin{aligned}
 u = & -\frac{Gm}{2M} \left(e^{-2\eta\sqrt{t}} \left[e^{\frac{4Mt}{Sc-1} + 2\eta\sqrt{t + \frac{4tMSc}{Sc-1}}} \operatorname{erfc} \left(\eta + \sqrt{t + \frac{4tMSc}{Sc-1}} \right) + e^{\frac{4Mt}{Sc-1} - 2\eta\sqrt{t + \frac{4tMSc}{Sc-1}}} \operatorname{erfc} \left(\eta - \sqrt{t + \frac{4tMSc}{Sc-1}} \right) - e^{2\eta(\sqrt{t+4Mt})} \cdot \operatorname{erfc}(\eta + \sqrt{t + 4Mt}) - e^{-2\eta(\sqrt{t+4Mt})} \cdot \operatorname{erfc}(\eta - \sqrt{t + 4Mt}) \right] \right. \\
 & - \frac{Gm\epsilon}{4M - \omega Sc i + \omega i} \left(e^{-(2\eta\sqrt{t})} \left[e^{\frac{4Mt}{Sc-1} + 2\eta\sqrt{t + \frac{4tMSc}{Sc-1}}} \cdot \operatorname{erfc} \left(\eta + \sqrt{t + \frac{4tMSc}{Sc-1}} \right) + e^{\frac{4Mt}{Sc-1} - 2\eta\sqrt{t + \frac{4tMSc}{Sc-1}}} \cdot \operatorname{erfc} \left(\eta - \sqrt{t + \frac{4tMSc}{Sc-1}} \right) - e^{t\omega i - 2\eta\sqrt{t(4M+1+\omega i)}} \cdot \operatorname{erfc} \left(\eta - \sqrt{t(4M+1+\omega i)} \right) - e^{t\omega i + 2\eta\sqrt{t(4M+1+\omega i)}} \cdot \operatorname{erfc} \left(\eta + \sqrt{t(4M+1+\omega i)} \right) \right] \right) \\
 & - \frac{2Gr}{4M - PrN} \left(e^{-(4Mt+t+2\eta\sqrt{t})} \left[e^{t\left(1 + \frac{Pr(4M-N)}{Pr-1}\right) + 2\eta\sqrt{t + \frac{Pr t(4M-N)}{Pr-1}}} \operatorname{erfc} \left(\eta + \sqrt{t + \frac{Pr t(4M-N)}{Pr-1}} \right) + e^{t\left(1 + \frac{Pr(4M-N)}{Pr-1}\right) - 2\eta\sqrt{t + \frac{Pr t(4M-N)}{Pr-1}}} \operatorname{erfc} \left(\eta - \sqrt{t + \frac{Pr t(4M-N)}{Pr-1}} \right) - e^{t(1+4M) + 2\eta\sqrt{t(1+4M)}} \operatorname{erfc} \left(\eta + \sqrt{t(1+4M)} \right) - e^{t(1+4M) - 2\eta\sqrt{t(1+4M)}} \operatorname{erfc} \left(\eta - \sqrt{t(1+4M)} \right) \right] \right) \\
 & - \frac{2Gr\epsilon}{4M - NPr - Pr\omega i + \omega i} \left(e^{-(4Mt+t+2\eta\sqrt{t})} \left[e^{t\left(1 + \frac{Pr(4M-N)}{Pr-1}\right) + 2\eta\sqrt{t + \frac{Pr t(4M-N)}{Pr-1}}} \operatorname{erfc} \left(\eta + \sqrt{t + \frac{Pr t(4M-N)}{Pr-1}} \right) + e^{t\left(1 + \frac{Pr(4M-N)}{Pr-1}\right) - 2\eta\sqrt{t + \frac{Pr t(4M-N)}{Pr-1}}} \operatorname{erfc} \left(\eta - \sqrt{t + \frac{Pr t(4M-N)}{Pr-1}} \right) - e^{t(1+4M+\omega i) + 2\eta\sqrt{t(1+4M+\omega i)}} \operatorname{erfc} \left(\eta + \sqrt{t(1+4M+\omega i)} \right) - e^{t(1+4M+\omega i) - 2\eta\sqrt{t(1+4M+\omega i)}} \operatorname{erfc} \left(\eta - \sqrt{t(1+4M+\omega i)} \right) \right] \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{Gm}{2M} \left(e^{-(2\eta Sc\sqrt{t})} \left[e^{\left(\frac{4Mt}{Sc-1}\right) + 2\eta\sqrt{tSc^2 + \frac{4MSct}{Sc-1}}} \cdot \operatorname{erfc} \left(\eta\sqrt{Sc} + \sqrt{tSc + \frac{4Mt}{Sc-1}} \right) + \right. \right. \\
 & e^{\left(\frac{4Mt}{Sc-1}\right) - 2\eta\sqrt{tSc^2 + \frac{4MSct}{Sc-1}}} \cdot \operatorname{erfc} \left(\eta\sqrt{Sc} - \sqrt{tSc + \frac{4Mt}{Sc-1}} \right) - e^{2\eta Sc\sqrt{t}} \cdot \operatorname{erfc}(\eta\sqrt{Sc} + \sqrt{tSc}) - \\
 & \left. \left. e^{-2\eta Sc\sqrt{t}} \cdot \operatorname{erfc}(\eta\sqrt{Sc} - \sqrt{tSc}) \right] \right) \\
 & + \frac{2Gme}{(4M - Sc\omega i + \omega i)} \left(e^{-(2\eta Sc\sqrt{t})} \left[e^{\left(tSc + \frac{4Mt}{Sc-1}\right) + 2\eta\sqrt{tSc^2 + \frac{4MSct}{Sc-1}}} \cdot \operatorname{erfc} \left(\eta\sqrt{Sc} + \sqrt{tSc + \frac{4Mt}{Sc-1}} \right) + \right. \right. \\
 & e^{\left(tSc + \frac{4Mt}{Sc-1}\right) - 2\eta\sqrt{tSc^2 + \frac{4MSct}{Sc-1}}} \cdot \operatorname{erfc} \left(\eta\sqrt{Sc} - \sqrt{tSc + \frac{4Mt}{Sc-1}} \right) - \\
 & \left. \left. e^{(\omega ti) + 2\eta\sqrt{Sc^2 t + Sc\omega ti}} \cdot \operatorname{erfc}(\eta\sqrt{Sc} + \sqrt{Sc t + \omega ti}) - e^{(\omega ti) - 2\eta\sqrt{Sc^2 t + Sc\omega ti}} \cdot \operatorname{erfc}(\eta\sqrt{Sc} - \right. \right. \\
 & \left. \left. \sqrt{Sc t + \omega ti}) \right] \right) \\
 & + \frac{2Gr}{4M - PrN} \left(e^{-(Pr t + N t + 2\eta Pr\sqrt{t})} \left[e^{\left(Pr t + \frac{t(4M-N)}{Pr-1}\right) + \left(2\eta\sqrt{tPr^2 + \frac{Pr t(4M-N)}{Pr-1}}\right)} \operatorname{erfc} \left(\eta\sqrt{Pr} + \right. \right. \\
 & \left. \left. \sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) + e^{\left(Pr t + \frac{t(4M-N)}{Pr-1}\right) - \left(2\eta\sqrt{tPr^2 + \frac{Pr t(4M-N)}{Pr-1}}\right)} \operatorname{erfc} \left(\eta\sqrt{Pr} - \sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) - \right. \\
 & \left. \left. e^{Pr t + tN + 2\eta\sqrt{Pr^2 t + PrNt}} \operatorname{erfc}(\eta\sqrt{Pr} + \sqrt{Pr t + Nt}) - e^{Pr t + tN - 2\eta\sqrt{Pr^2 t + PrNt}} \operatorname{erfc}(\eta\sqrt{Pr} - \right. \right. \\
 & \left. \left. \sqrt{Pr t + Nt}) \right] \right) \\
 & + \frac{2Gre}{4M - PrN - Pr\omega i + \omega i} \left(e^{-(Pr t + N t + 2\eta Pr\sqrt{t})} \right)
 \end{aligned}$$

$$\left[e^{\left(\frac{Pr t + \frac{t(4M-N)}{Pr-1}}{Pr-1}\right) + \left(2\eta\sqrt{tPr^2 + \frac{Pr t(4M-N)}{Pr-1}}\right)} \operatorname{erfc}\left(\eta\sqrt{Pr} + \sqrt{Pr t + \frac{t(4M-N)}{Pr-1}}\right) + e^{\left(\frac{Pr t + \frac{t(4M-N)}{Pr-1}}{Pr-1}\right) - \left(2\eta\sqrt{tPr^2 + \frac{Pr t(4M-N)}{Pr-1}}\right)} \operatorname{erfc}\left(\eta\sqrt{Pr} - \sqrt{Pr t + \frac{t(4M-N)}{Pr-1}}\right) - e^{t(Pr+N+\omega i) + 2\eta\sqrt{t(Pr^2+NPr+Pr\omega i)}} \cdot \operatorname{erfc}\left(\eta\sqrt{Pr} + \sqrt{t(Pr+N+\omega i)}\right) - e^{t(Pr+N+\omega i) - 2\eta\sqrt{t(Pr^2+NPr+Pr\omega i)}} \cdot \operatorname{erfc}\left(\eta\sqrt{Pr} - \sqrt{t(Pr+N+\omega i)}\right) \right] \dots (19)$$

$$T = \frac{e^{-2\eta Pr\sqrt{t}}}{2} \left[e^{-2\eta\sqrt{tPr^2+tNPr}} \cdot \operatorname{erfc}\left(\eta\sqrt{Pr} - \sqrt{t(Pr+N)}\right) + e^{2\eta\sqrt{tPr^2+tNPr}} \cdot \operatorname{erfc}\left(\eta\sqrt{Pr} + \sqrt{t(Pr+N)}\right) + \epsilon \left(e^{t\omega i - 2\eta\sqrt{tPr^2+tNPr+tPr\omega i}} \cdot \operatorname{erfc}\left(\eta\sqrt{Pr} - \sqrt{t(Pr+N+\omega i)}\right) + e^{t\omega i + 2\eta\sqrt{tPr^2+tNPr+tPr\omega i}} \cdot \operatorname{erfc}\left(\eta\sqrt{Pr} + \sqrt{t(Pr+N+\omega i)}\right) \right) \right] \dots (20)$$

$$C = \frac{e^{-2\eta Sc\sqrt{t}}}{2} \left[e^{-2\eta Sc\sqrt{t}} \cdot \operatorname{erfc}\left(\eta\sqrt{Sc} - \sqrt{tSc}\right) + e^{2\eta Sc\sqrt{t}} \cdot \operatorname{erfc}\left(\eta\sqrt{Sc} + \sqrt{tSc}\right) + \epsilon \left(e^{t\omega i - 2\eta\sqrt{tSc^2+tSc\omega i}} \cdot \operatorname{erfc}\left(\eta\sqrt{Sc} - \sqrt{t(Sc+\omega i)}\right) + e^{t\omega i + 2\eta\sqrt{tSc^2+tSc\omega i}} \cdot \operatorname{erfc}\left(\eta\sqrt{Sc} + \sqrt{t(Sc+\omega i)}\right) \right) \right] \dots (21)$$

Where, $\eta = \frac{y}{4\sqrt{t}}$

The non-dimensional shearing stress at the wall from the equations (19) is given by

$$\begin{aligned} \tau = \left(\frac{\partial u}{\partial \eta}\right)_{\eta=0} &= \frac{Gm\sqrt{t}}{M} \left[e^{\frac{4Mt}{Sc-1}} \left(\operatorname{erfc}\left(\sqrt{t + \frac{4MtSc}{Sc-1}}\right) + \operatorname{erfc}\left(-\sqrt{t + \frac{4MtSc}{Sc-1}}\right) \right) - \right. \\ &\left. \left(\operatorname{erfc}(\sqrt{t + 4Mt}) + \operatorname{erfc}(-\sqrt{t + 4Mt}) \right) \right] - \\ &\frac{Gm}{2M} \left[2\sqrt{t + \frac{4MtSc}{Sc-1}} e^{\frac{4Mt}{Sc-1}} \left(\operatorname{erfc}\left(\sqrt{t + \frac{4MtSc}{Sc-1}}\right) - \operatorname{erfc}\left(-\sqrt{t + \frac{4MtSc}{Sc-1}}\right) \right) - \right. \\ &\left. 2\sqrt{t + 4Mt} \left(\operatorname{erfc}(\sqrt{t + 4Mt}) - \operatorname{erfc}(-\sqrt{t + 4Mt}) \right) \right] + \\ &\frac{4Gme\sqrt{t}}{4M - \omega Sci + \omega i} e^{-(1+4M)\epsilon} \left[e^{\frac{4MSct}{Sc-1}} \left(\operatorname{erfc}\left(\sqrt{t + \frac{4MtSc}{Sc-1}}\right) + \operatorname{erfc}\left(-\sqrt{t + \frac{4MtSc}{Sc-1}}\right) \right) - \right. \\ &\left. e^{t(4M+1+\omega i)} \left(\operatorname{erfc}(\sqrt{t + 4Mt + \omega ti}) + \operatorname{erfc}(-\sqrt{t + 4Mt + \omega ti}) \right) \right] \\ &- \frac{2Gme}{4M - \omega Sci + \omega i} e^{-(1+4M)\epsilon} \left[2\sqrt{t + \frac{4MtSc}{Sc-1}} e^{\frac{4MSct}{Sc-1}} \left(\operatorname{erfc}\left(\sqrt{t + \frac{4MtSc}{Sc-1}}\right) - \right. \right. \\ &\left. \left. \operatorname{erfc}\left(-\sqrt{t + \frac{4MtSc}{Sc-1}}\right) \right) - 2\sqrt{t + 4Mt + \omega ti} e^{t(4M+1+\omega i)} \left(\operatorname{erfc}(\sqrt{t + 4Mt + \omega ti}) - \right. \right. \\ &\left. \left. \operatorname{erfc}(-\sqrt{t + 4Mt + \omega ti}) \right) \right] + \\ &\frac{4Gr\sqrt{t}}{4M - PrN} \left[e^{\frac{t(4M-N)}{Pr-1}} \left(\operatorname{erfc}\left(\sqrt{t + \frac{Prt(4M-N)}{Pr-1}}\right) + \operatorname{erfc}\left(-\sqrt{t + \frac{Prt(4M-N)}{Pr-1}}\right) \right) - \right. \\ &\left. \left(\operatorname{erfc}(\sqrt{t + 4Mt}) + \operatorname{erfc}(-\sqrt{t + 4Mt}) \right) \right] - \\ &\frac{2Gr}{4M - PrN} \left[2\sqrt{t + \frac{Prt(4M-N)}{Pr-1}} e^{\frac{t(4M-N)}{Pr-1}} \left(\operatorname{erfc}\left(\sqrt{t + \frac{Prt(4M-N)}{Pr-1}}\right) - \right. \right. \\ &\left. \left. \operatorname{erfc}\left(-\sqrt{t + \frac{Prt(4M-N)}{Pr-1}}\right) \right) - 2\sqrt{t + 4Mt} \left(\left(\operatorname{erfc}(\sqrt{t + 4Mt}) - \operatorname{erfc}(-\sqrt{t + 4Mt}) \right) \right) \right] \end{aligned}$$

$$\begin{aligned}
 & + \frac{4Gre\sqrt{t}}{4M - NPr - Pr\omega i + \omega i} \left[e^{\frac{t(4M - PrN)}{Pr - 1}} \left(\operatorname{erfc} \left(\sqrt{t + \frac{Pr t(4M - N)}{Pr - 1}} \right) + \operatorname{erfc} \left(-\sqrt{t + \frac{Pr t(4M - N)}{Pr - 1}} \right) \right) - \right. \\
 & \left. e^{t\omega i} \left(\operatorname{erfc} \left(\sqrt{t(1 + 4M + \omega i)} \right) + \operatorname{erfc} \left(-\sqrt{t(1 + 4M + \omega i)} \right) \right) \right] - \\
 & \frac{2Gre}{4M - NPr - Pr\omega i + \omega i} \left[2\sqrt{t + \frac{Pr t(4M - N)}{Pr - 1}} e^{\frac{t(4M - PrN)}{Pr - 1}} \left(\operatorname{erfc} \left(\sqrt{t + \frac{Pr t(4M - N)}{Pr - 1}} \right) - \right. \right. \\
 & \left. \left. \operatorname{erfc} \left(-\sqrt{t + \frac{Pr t(4M - N)}{Pr - 1}} \right) \right) - 2\sqrt{t(1 + 4M + \omega i)} e^{t\omega i} \left(\operatorname{erfc} \left(\sqrt{t(1 + 4M + \omega i)} \right) - \right. \right. \\
 & \left. \left. \operatorname{erfc} \left(-\sqrt{t(1 + 4M + \omega i)} \right) \right) \right] - \\
 & \frac{4GmSc\sqrt{t}}{Sc^2 - Sc + 4M} \left[e^{\frac{4Mt}{Sc - 1}} \left(\operatorname{erfc} \left(\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) + \operatorname{erfc} \left(-\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) \right) - \left(\operatorname{erfc}(\sqrt{Sct}) + \right. \right. \\
 & \left. \left. \operatorname{erfc}(-\sqrt{Sct}) \right) \right] + \\
 & \frac{Gm}{2M} \left[2\sqrt{tSc^2 + \frac{4ScMt}{Sc - 1}} e^{\frac{4Mt}{Sc - 1}} \left(\operatorname{erfc} \left(\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) - \operatorname{erfc} \left(-\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) \right) - \right. \\
 & \left. 2Sc\sqrt{t} \left(\operatorname{erfc}(\sqrt{Sct}) - \operatorname{erfc}(-\sqrt{Sct}) \right) \right] - \frac{4GmSc\sqrt{t}}{4M - Sc\omega i + \omega i} \left[e^{\frac{4Mt}{Sc - 1}} \operatorname{erfc} \left(\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) + \right. \\
 & \left. \operatorname{erfc} \left(-\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) - e^{t\omega i} \left(\operatorname{erfc}(\sqrt{Sct + \omega ti}) + \operatorname{erfc}(-\sqrt{Sct + \omega ti}) \right) \right] \\
 & + \\
 & \frac{2Gme}{4M - Sc\omega i + \omega i} \left[2\sqrt{tSc^2 + \frac{4ScMt}{Sc - 1}} e^{\frac{4Mt}{Sc - 1}} \left(\operatorname{erfc} \left(\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) - \operatorname{erfc} \left(-\sqrt{tSc + \frac{4Mt}{Sc - 1}} \right) \right) - \right. \\
 & \left. 2\sqrt{Sc^2 t + Sc\omega ti} \cdot e^{t\omega i} \left(\operatorname{erfc}(\sqrt{Sct + \omega ti}) - \operatorname{erfc}(-\sqrt{Sct + \omega ti}) \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{4GrPr\sqrt{t}}{4M-PrN} \left[e^{\frac{t(4M-PrN)}{Pr-1}} \left(\operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) + \operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) \right) - \right. \\
 & \left. \left(\operatorname{erfc}(\sqrt{Pr t + Nt}) + \operatorname{erfc}(-\sqrt{Pr t + Nt}) \right) \right] + \\
 & \frac{2Gr}{4M-PrN} \left[2\sqrt{tPr^2 + \frac{Pr t(4M-N)}{Pr-1}} e^{\frac{t(4M-PrN)}{Pr-1}} \left(\operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) - \right. \right. \\
 & \left. \left. \operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) \right) - \right. \\
 & \left. 2\sqrt{Pr^2 t + NPr t} \left(\operatorname{erfc}(\sqrt{Pr t + Nt}) - \operatorname{erfc}(-\sqrt{Pr t + Nt}) \right) \right] - \\
 & \frac{4GrePr\sqrt{t}}{4M-PrN-Pr\omega i + \omega i} \left[e^{\frac{t(4M-PrN)}{Pr-1}} \left(\operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) + \operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) \right) - \right. \\
 & \left. e^{t\omega i} \left(\operatorname{erfc}(\sqrt{Pr t + Nt + \omega t i}) + \operatorname{erfc}(-\sqrt{Pr t + Nt + \omega t i}) \right) \right] + \\
 & \frac{2Gre}{4M-PrN-Pr\omega i + \omega i} \left[2\sqrt{tPr^2 + \frac{Pr t(4M-N)}{Pr-1}} e^{\frac{t(4M-PrN)}{Pr-1}} \left(\operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) - \right. \right. \\
 & \left. \left. \operatorname{erfc} \left(\sqrt{Pr t + \frac{t(4M-N)}{Pr-1}} \right) \right) - 2\sqrt{Pr^2 t + NPr t + \omega t i} e^{t\omega i} \left(\operatorname{erfc}(\sqrt{Pr t + Nt + \omega t i}) - \right. \right. \\
 & \left. \left. \operatorname{erfc}(-\sqrt{Pr t + Nt + \omega t i}) \right) \right]
 \end{aligned}
 \tag{22}$$

The dimensional rate of heat transfer / Nusselt Number,

$$\begin{aligned}
 Nu &= -\left(\frac{\partial T}{\partial \eta}\right)_{\eta=0} = Pr\sqrt{t} \left[\operatorname{erfc}(-\sqrt{t(Pr+N)}) + \operatorname{erfc}(\sqrt{t(Pr+N)}) + e^{t\omega i} \left(\operatorname{erfc}(-\sqrt{t(Pr+N+\omega i)}) + \right. \right. \\
 & \left. \left. \operatorname{erfc}(\sqrt{t(Pr+N+\omega i)}) \right) \right] - \\
 & \left[\sqrt{tPr^2 + tPrN} \left(\operatorname{erfc}(\sqrt{t(Pr+N)}) - \operatorname{erfc}(-\sqrt{t(Pr+N)}) \right) - \frac{4\sqrt{Pr}}{\sqrt{\pi}} e^{-t(Pr+N)} + \right. \\
 & \left. \sqrt{tPr^2 + tPrN + tPr\omega i} \cdot e^{t\omega i} \left(\operatorname{erfc}(\sqrt{t(Pr+N+\omega i)}) - \operatorname{erfc}(-\sqrt{t(Pr+N+\omega i)}) \right) \right]
 \end{aligned}
 \tag{23}$$

The dimensionless rate of mass transfer/ Sherwood Number,

$$Sh = - \left(\frac{\partial C}{\partial \eta} \right)_{\eta=0} = Sc\sqrt{t} \left[\operatorname{erfc}(-\sqrt{tSc}) + \operatorname{erfc}(\sqrt{tSc}) + e^{ti\omega} \left(\operatorname{erfc}(-\sqrt{t(Sc + \omega i)}) + \operatorname{erfc}(\sqrt{t(Sc + \omega i)}) \right) \right] - \left[Sc\sqrt{t} \left(-\operatorname{erfc}(-\sqrt{tSc}) + \operatorname{erfc}(\sqrt{tSc}) \right) - \frac{2\sqrt{Sc}}{\sqrt{\pi}} e^{-tSc} + e^{ti\omega} \sqrt{tSc^2 + tSc\omega i} \left(-\operatorname{erfc}(-\sqrt{t(Sc + \omega i)}) + \operatorname{erfc}(\sqrt{t(Sc + \omega i)}) \right) - \frac{2\sqrt{Sc}}{\sqrt{\pi}} e^{-t(Sc + \omega i)} \right] \dots (24)$$

In order to get the physical insight into the problem, the numerical value of u, τ and other Physical quantities have been calculated from the above equations. At the time of evaluation of the expressions, it is observed that the argument of the error function is complex and hence, it is separated into real and imaginary parts by using the following formula and then the complimentary error function up to twenty terms is calculated.

$$\operatorname{erfc}(a + ib) = \operatorname{erf}(a) + \frac{\exp(-a^2)}{2a\pi} [1 - \cos(2ab) + i \sin(2ab)] + \frac{2 \exp(-n^2/4)}{\pi} \sum_{n=1}^{\infty} [f_n(a,b) + i g_n(a,b)] + \varepsilon(a,b) \dots (25)$$

Where,

$$f_n = 2a - 2a \cosh(nb) \cos(2ab) + n \sinh(nb) \sin(2ab),$$

$$g_n = 2a \cosh(nb) \sin(2ab) + n \sinh(nb) \cos(2ab)$$

and $|\varepsilon(a,b)| \approx 10^{-16} |\operatorname{erf}(a + ib)|$

Discussion

In this paper, we have studied the effect of radiation on unsteady MHD free convective flow past an infinite vertical plate with heat, mass transfer and constant suction. The effect of parameters Gr, Gm, M, N, t, Pr, ω, Sc, on flow characteristics has been presented in the (Fig.1-9).

Velocity profile: The mean velocity profiles are depicted in (Fig.1-2). Fig.1 shows the effect of the parameters Gr, Gm & M on velocity at any point of the fluid, when N=1, Pr=2 & t=1. It is observed that the velocity decreases with both the parameters Gr & Gm very near the plate and then increases away from the plate. But, the velocity increases for




Fig 1: Effect of Gr, Gm and M on velocity profile when N=1, Pr=2 and t=1




Fig 2: Effect of N, Pr and t on velocity profile when Gr=5, Gm=5 and M=5




Fig 3: Effect of N, Pr, ω and t on heat profile when $\epsilon = 0.02$

the parameter M throughout diminishing to zero at a small distance from the plate. Fig. 2 Shows the effect of the parameters N, Pr & t on velocity at any point of the fluid, when Gr=5, Gm=5 and M=5. It is observed that the velocity increases with both the parameters Pr & N very near the plate and decreases away from the plate. But the velocity increases for the parameter t throughout diminishing to zero at a small distance from the plate.




Fig 4: Effect of Sc, ω and t on mass concentration profile when $\epsilon = 0.02$




Fig 5: Effect of Pr, ω and N on Nusselt Number when $\epsilon = 0.02$

Temperature profile: The temperature profile is depicted in (Fig. 2). Fig.3 shows the effect of the parameters Pr, N, ω and t on heat at any point of the fluid, when $\epsilon=0.02$. It is observed that the temperature falls with increase of the parameters Prandtl number (Pr), Oscillating frequency (ω) and time (t).

Concentration profile: The concentration profile is depicted in Fig. 4. Fig. 4 shows the effect of the parameters Sc, ω and t on concentration at any point of the fluid, when $\epsilon=0.02$. It is noticed that




Fig 6: Effect of Sc and ω on Sherwood Number when $\epsilon = 0.02$




Fig 8: Effect of ω and M on shearing stress when $Gr=5$, $Sc=2$, $Pr=2$, $Gm=5$, $\epsilon = 0.02$ and $N=1$




Fig 7: Effect of Gr and N on shearing stress when $M=1$, $Sc=2$, $Pr=2$, $Gm=5$, $\epsilon = 0.02$ and $\omega=1$




Fig 9: Effect of Sc , Pr and Gm on shearing stress when $Gr=5$, $\omega=1$, $M=1$, $\epsilon = 0.02$ and $N=1$

the Concentration decreases with increase of all the parameters Schmidt number (Sc), oscillating frequency (ω) and time (t).

Nusselt Number: The Nusselt number is depicted in (Fig. 5). Fig.5 shows the effect of the parameters Pr , ω and N on Nusselt number at any point of the fluid, when $\epsilon=0.02$. It is noticed that the Nusselt number increases with the increase of the parameter Prandtl number (Pr) and Radiation parameter (N) but decreases with the Oscillating frequency (ω).

Sherwood Number: The Sherwood number is depicted in Fig. 6. Fig. 6 shows the effect of the parameters Sc and ω on Sherwood number at any point of the fluid, when $\epsilon=0.02$. It is noticed that the Sherwood number increases with the increase of the parameter Schmidt number (Sc), but decreases with the Oscillating frequency (ω).

Shearing Stress: The Shearing Stress is depicted in (Fig 7-9). (Fig. 7) shows the effect of the parameters Gr and N on Shearing Stress at any point of the fluid, when $M=1$, $Sc=2$, $Pr=2$, $Gm=5$,

$\epsilon=0.02$ and $\omega=1$. It is noticed that the Shearing Stress increases with the increase of the parameter Grashof number (Gr), but decreases with the Radiation parameter (N). Fig. 8 shows the effect of the parameters ω and M on Shearing Stress at any point of the fluid, when $Gr=5$, $Sc=2$, $Pr=2$, $Gm=5$, $\epsilon=0.02$ and $N=1$. It is noticed that the Shearing Stress decreases with decrease of the parameter Oscillating frequency (ω), but increases with the Magnetic number(M). Figure-(9) shows effect of the parameters Sc, Pr and Gm on Shearing Stress at any point of the fluid, when $Gr=5$, $\omega=1$, $N=1$, $\epsilon=0.02$ and $M=1$. Shearing Stress has no effect. For the parameters Schmidt number (Sc) & modified Grashof number (Gm).

Sherwood Number: The Sherwood number is depicted in (Fig. 6). Fig. 6 shows the effect of the parameters Sc and ω on Sherwood number at any point of the fluid, when $\epsilon=0.02$. It is noticed that the Sherwood number increases with the increase of the parameter Schmidt number (Sc), but decreases with the Oscillating frequency (ω).

Shearing Stress: The Shearing Stress is depicted in Fig.7-9. Fig. 7 shows the effect of the parameters Gr and N on Shearing Stress at any point of the fluid, when $M=1$, $Sc=2$, $Pr=2$, $Gm=5$, $\epsilon=0.02$ and $\omega=1$. It is noticed that the Shearing Stress increases with the increase of the parameter Grashof number (Gr), but decreases with the Radiation parameter (N). Fig. 8 shows the effect of the parameters ω and M on Shearing Stress at any point of the fluid, when $Gr=5$, $Sc=2$, $Pr=2$, $Gm=5$, $\epsilon=0.02$ and $N=1$. It is noticed that the Shearing Stress decreases with decrease of the parameter Oscillating frequency (ω), but increases with the Magnetic number (M). Fig.9 shows effect of the parameters Sc, Pr and Gm on Shearing Stress at any point of the fluid, when $Gr=5$, $\omega=1$, $N=1$, $\epsilon=0.02$ and $M=1$. Shearing Stress has no effect. For the parameters Schmidt number (Sc) & modified Grashof number (Gm).

References

- Acharya M, Dash GC & Singh LP (2000) Magnetic Field Effects on the Free Convection and Mass Transfer Flow Through Porous Medium with Constant Suction and Constant Heat Flux, *Indian J. Pure Appl. Math.* 31(1), 1-18.
- Ahmed Y. Ghaly: Department of Mathematics, Faculty of Education, Ain Shams University, Roxy, year ppno 93-103.
- Carnahan P, Luther HA & Wilkes J O (1969) Applied Numerical Methods, John Wiley & Sons, USA.
- Chamkha AJ (2004) Unsteady MHD Convective Heat and Mass Transfer Past A Semi-Infinite Vertical Permeable Moving Plate with Heat Absorption, *Int. J. Engg. Sci.* 42, 217-230.
- Chen CK & Char MI (1998) Heat transfer of a continuous, stretching surface with suction or blowing, *J. Math. Anal. Appl.* 135(2), 568-580.
- Danberg JE & Fansber KS (1976). Non similar solution for the flow in the boundary layer past a stretched wall, *Quart. Appl. Math.* 34, 305-311.
- Elbasheshy, EMA (1998) Heat transfer over a stretching surface with variable surface heat flux, *J. Phys. D: Appl. Phys.* 31, 1951-1954.
- Erickson LE, Fan LT & Fox VG (1966) Heat and mass transfer on a moving continuous flat plate with suction or injection, *Indust. Eng. Chem.* 5, 19-25.
- Gupta PS & Gupta AS (1977) Heat and mass transfer on a stretching sheet with suction or blowing, *Canadian J. Chem. Engrg.* 55, 744-746.
- Helmy KA (1998) MHD Unsteady Free Convection Flow past a Vertical Porous Plate, *ZAMM*, 78(4), 225-270.
- Muthucumaraswamy R & Kumar GS (2004) Heat and Mass Transfer Effects on Moving Vertical Plate in The Presence of Thermal Radiation, *Theoret. Appl. Mech.* 31(1), 35-46.
- Rohsenow WM, Hartnett JP, & Cho YI (1998) (eds.), *Handbook of Heat Transfer*, 3rd ed., McGraw-Hill, New York, 1998.
- Sahoo PK, Datta N & Biswal S (2003) Magnetohydrodynamic Unsteady Free Convection Flow Past an Infinite Vertical Plate With Constant Suction and Heat Sink, *Indian J. Pure Appl. Math.*, 34(1), 145-155.
- Sakiadis BC (1961) Boundary layer behaviour on continuous solid surfaces: I. The boundary layer



equations for two-dimensional and axisymmetric flow, *AIChE J.* 7, 26–28.

15. Singh AK, Singh AK & Singh N P (2003) Heat and Mass Transfer In MHD Flow of a Viscous Fluid Past a Vertical Plate Under Oscillatory Suction Velocity, *Indian. J. Pure Appl. Math.*, 34(3), 429-442.
16. Takhar HS, Gorla RSR, & Soundalgekar VM (1996) Radiation effects on MHD free convection flow of a gas past a semi-infinite vertical plate, *Internat. J. Numer. Methods Heat Fluid Flow*, 6(2), 77–83.
17. Vajravelu K & Hadjinicolaou, A (1997) Convective heat transfer in an electrically 10.conducting fluid at a stretching surface with uniform free stream, *Internat. J. Engrg. Sci.* 35, 12-13, 1237–1244.
18. Young JK (2000) Unsteady MHD Convective Heat Transfer Past A Semi-Infinite Vertical Porous Moving Plate with Variable Suction, *Int. J. Engg. Sci.* 38, 833-845.