

# Antioxidant Activity of Geraniol on n-Nitrosodiethylamine-induced Hepatocarcinogenesis in Wistar Albino Rats

Fouzia Banu<sup>1</sup>, Nausheen Dawood<sup>1</sup>, Yamini Sudha Lakshmi<sup>2\*</sup>, S. Gopalakrishnan<sup>3</sup> and V. Brinda<sup>4</sup>

<sup>1</sup>Department of Biochemistry, JBAS College for Women, Chennai, India

<sup>2</sup>Prof. Dhanapalan College of Arts and Science, Chennai, India; yasula2000@yahoo.com

<sup>3</sup>PNG University of Technology

<sup>4</sup>CLRI, Chennai

## Abstract

The chemopreventive effect of Geraniol was evaluated in N-nitrosodiethylamine-induced experimental liver tumor in Wistar albino rats. In this study the animals were divided into four groups with 6 rats each and received treatments for 4 weeks. Group I served as control (untreated), Group II treated with N-nitrosodiethylamine (DEN) (200 mg/kg, i.p.) and Phenobarbital (PB) (0.05%) in drinking water; Group III received Geraniol (250 mg/kg b.wt) in addition to DEN and Group IV treated with Geraniol only for four weeks. The administration of Geraniol in rats with DEN induced HCC rats (Group III) shows a significantly reduced level of liver serum marker (aspartate aminotransferase, alanine aminotransferase, acid phosphatase, alkaline phosphatase, lactate dehydrogenase, gammaglutamyl transpeptidase, alpha fetoprotein and 5' nucleotidase). These results are evident that Geraniol possesses protective effect against N-Nitrosodiethylamine (DEN)-induced Hepatocellular Carcinogenesis (HCC).

**Keywords:** DEN, Carcinogenesis, HCC, Geraniol, Marker Enzymes

## 1. Introduction

Oxygen free radicals are generated during various metabolic processes and the antioxidant defense mechanism of the body keeps a check on the excess production of free radicals generated in the system<sup>1,2</sup>. The imbalance between the exposure to oxidants and the endogenous antioxidants results in 'oxidative stress' which is found to play an important role in various pathological conditions including asthma, Acute Respiratory Distress Syndrome (ARDS), Chronic Obstructive Pulmonary Disease (COPD), idiopathic pulmonary fibrosis, cardiovascular dysfunction, neurodegenerative diseases, immunosuppression, cancer, premature ageing etc.<sup>3,4</sup>.

The liver is a vital organ where free radicals are largely generated and a large number of plants have claimed to have hepatoprotective activity<sup>5,6</sup>.

Diethyl Nitrosamine (DEN) is one of the most important environmental carcinogens in N-nitrosamines class, which primarily induces tumours of the liver<sup>7</sup>. DEN is considered to be a genotoxic carcinogen<sup>8</sup>. It is often assumed that DEN initiates and propagates tumor development primarily by inducing DNA alterations that lead to mutations<sup>9</sup>.

Herbs have been used as food and for medicinal purposes for centuries. Research interest has focused on various herbs that possess hypolipidemic, antiplatelet, antitumor or immune-stimulating properties that may be useful adjuncts in helping reduce the risk of cardiovascular disease and cancer. In different herbs, a wide variety of active phytochemicals including the flavonoids, terpenoids, lignins, sulphides, polyphenolics, carotenoids, coumarines, saponins, plant sterols, curcumins and phthalides have been identified<sup>10</sup>.

\*Author for correspondence

Natural essential oils contain monoterpenes as one of their major constituents. The oil accumulates in glands found in leaves, petals and trichomes, and geraniol is an acyclic monoterpene alcohol that is found in the essential oils of *Cymbopogon citratus*, lemongrass<sup>11</sup>, *Zingiber officinale*, ginger<sup>12</sup> and *Ocimum basilicum*, basil<sup>13</sup>. It is also emitted from flowers of many plant species with an intense rose scent. In the camphor tree, *Cinnamomum tenuipile* the leaves accumulate almost 86–98% geraniol in their essential oil<sup>14</sup>.

In recent years, the general public has shown an increased interest in the use of herbal medicines in preference to synthetic drugs. This is based on the belief that natural products are intrinsically less dangerous and can be obtained at a lower cost<sup>15</sup>.

Chemoprevention, a novel and appealing strategy, deals with the inhibition, reversal or suppression of carcinogenesis by the use of natural or synthetic agents<sup>16</sup>.

Phytochemicals such as alkaloids, monoterpenes, flavanoids, isoflavones, saponins etc. not be considered nutritive compounds, they are known to possess protective activity against diseases. These chemicals are known to be produced by plants for their own protection but their pharmacological effects are of great interest and are being explored extensively. Geraniol, an important constituent of essential oil of ginger, lemon, lime, lavender, nutmeg, orange, rose etc is an acyclic monoterpene and is the main component of oil-of-rose and palmarosa oil. It is used in perfumes due to its rose like fragrance. Geraniol, a colourless liquid, is an acyclic terpene alcohol with a flowery rose-like odour. It has been assessed by the Joint FAO/WHO Expert Committee on Food Additives as safe at current levels of intake as flavoring agents. In USA the daily intake is estimated at 5.2 lg/kg bw/day while in Europe it is estimated 11 lg/kg bw/day<sup>17</sup>. Geraniol is reported to exert anti-tumor activity against various cancer cells both in vitro and in vivo<sup>18–20</sup>.

## 2. Materials and Methods

### 2.1 Materials

#### 2.1.1 Animals

Adult male Wistar albino rats were obtained from the King's Institute–Guindy, Chennai and housed in clean polypropylene cages. Animals were maintained according to the principle and guidelines of the committee (No. 945/C/01/04/06/CPCSEA) under the supervision of Animal

Ethical Committee. The animals were fed on commercial pelleted feed procured from Poultry Research Station, Chennai and water was provided *ad libitum*.

#### 2.1.2 Chemicals

Geraniol (98% pure), Diethylnitrosamine (DEN), Phenobarbital (PB) and Reduced glutathione were purchased from Sigma Chemical Company, St. Louis, MO, USA.

## 2.2 Methods

### 2.2.1 Experimental Design

Four groups with six animals each were taken for experiment.

- Group I : Control rats (untreated)
- Group II : Rats administered with Diethylnitrosamine (DEN) (200 mg/kg body weight) intraperitoneally and Phenobarbital (PB) (0.05%) in drinking water.
- Group III : Hepatocellular Carcinoma (HCC) bearing rats administered orally with Geraniol (250 mg/kg body weight).
- Group IV : Rats administered orally only with Geraniol (250 mg/kg body weight).

Group I rats fed with pellet feed and water served as normal control rats. Group II rats were treated with DEN by a single intraperitoneal injection (200 mg/kg body weight). After two weeks, the carcinogenic effect was promoted by Phenobarbital, the promoter (0.05% PB). The Promoter was supplemented to the animals through drinking water up to 14 successive weeks<sup>21</sup>. Group III animals were also treated with DEN by intraperitoneal injection (200 mg/kg body weight). After two weeks Phenobarbital was supplemented through drinking water for 14 successive weeks. At the beginning of 15<sup>th</sup> week, the cancer induced rats were treated with geraniol (250 mg/kg bodyweight) dissolved in corn oil (2.5 ml/kg body weight) orally for eight consecutive weeks. Group IV animals were treated only with Geraniol (250 mg/kg body weight) for 8 consecutive weeks to study the cytotoxicity (if any) induced by Geraniol.

At the end of the experiment, by cardiac puncture blood samples were collected and allowed to clot. From the clotted sample serum was separated by centrifugation at 3000 g for 10 min and stored at –70 °C for biochemical assays such as Aspartate Aminotransferase (AST),

Alanine Aminotransferase (ALT), Acid Phosphatase (ACP), Alkaline Phosphatase (ALP), Lactate Dehydrogenase (LDH), Gamma Glutamyl Transpeptidase (GTT), Alpha Feto-Protein (AFP) and 5'Nucleotidase levels.

## 2.3 Supernatant Preparation

The liver samples were harvested from the experimental animals and weighted and perfused immediately with ice-cold saline (0.85%, w/v NaCl), and homogenized with chilled phosphate buffer (0.1M, pH 7.4) containing KCl (1.17%, w/v). The homogenate was then centrifuged (800 g, 5 min, 4 °C) for removing the cell debris. The supernatant was collected and centrifuged at 10,000 g for 20 min at 4° C. Then the resulting supernatant is then assayed for CAT, SOD, and GPx activities.

## 2.4 Determination of the Extract Effect on Lipid Peroxidation in Liver

Lipid peroxidation was estimated by Thiobarbituric Acid (TBA) reaction with Malondialdehyde (MDA) as reported earlier<sup>22</sup>. Lipid peroxidation was expressed as nanomoles of MDA per milligram of protein.

## 2.5 Determination of the Extract Effect on Antioxidant Enzyme Activities in the Liver

### 2.5.1 Superoxide Dismutase Activity

The SOD activity was measured according to the method used by Marklund and Marklund<sup>23</sup>. The enzyme activity was expressed as units/mg protein.

### 2.5.2 Catalase Activity

The protein content of the supernatant was determined using the method with copper sulphate as reported earlier<sup>24</sup>.

The enzyme activity was calculated as nano mole of H<sub>2</sub>O<sub>2</sub> consumed/min/mg protein.

### 2.5.3 Glutathione Peroxidase Activity

To estimate the GPx activity, the reaction mixture consisted of 1.65 mL phosphate buffer (0.1 M, pH 7.4), 0.1 mL EDTA (0.5 mM), 0.05 mL oxidized glutathione (1 mM), 0.1 mL NADPH (0.1 mM), and 0.1 mL supernatant in a total volume of 2 mL. The disappearance of NADPH at 340 nm was recorded at 25 °C. The enzyme activity was calculated as nano mol of NADPH oxidized/min/mg protein using molar extinction coefficient.

## 3. Results

The activities of enzymatic antioxidants (SOD, CAT and GPx) level in the liver hemogenate of the experimental animals in each group are shown in Table 1.

The activities of SOD, CAT and GPx level were significantly decreased in hemogenate of tumor bearing animals (Group II) as compared to control animals. Geraniol administrated rats showed significant increase in the enzymatic antioxidants levels (Group III and Group IV).

The activities of enzymatic antioxidants (SOD, CAT and GPx) level in liver of the experimental animals in each group are shown in Table 2.

The activities of SOD, CAT and GPx level were significantly decreased in liver tissue of tumor bearing animals (Group II) as compared to control animals. Geraniol administrated rats showed significant increase in the enzymatic antioxidants levels (Group III and Group IV).

The activities of non-enzymatic antioxidants (GSH, Vit C and Vit E) level in haemolysate of the experimental animals in each group are shown in Table 3.

**Table 1.** Effect of geraniol on the enzymatic antioxidants in liver hemogenate of control and experimental animals

Parameters	Superoxide dismutase (SOD)	Catalase (CAT)	Glutathione peroxidase (GPx)
Group I (Control)	3.72 ± 0.08	65.92 ± 4.23	6.52 ± 0.25
Group II (DEN)	2.34 ± 0.33 <sup>****</sup>	42.44 ± 2.42 <sup>****</sup>	3.81 ± 0.32 <sup>a****</sup>
Group III (DEN + Geraniol)	3.38 ± 0.03 <sup>b****</sup>	57.25 ± 4.49 <sup>b****</sup>	4.40 ± 0.33 <sup>b**</sup>
Group IV (Geraniol)	3.59 ± 0.19 <sup>aNS</sup>	64.68 ± 2.76 <sup>aNS</sup>	6.42 ± 0.28 <sup>aNS</sup>

Values represent mean ± SD for 6 rats in each group.

Units: SOD – units /mg protein

CAT – μ moles of H<sub>2</sub>O<sub>2</sub> utilized/min/mg protein.

GPx – μ moles of GSH oxidised /min/mg protein.

\*\*\* p<0.001; \*\* p<0.01; \* p<0.05

<sup>NS</sup> Non-significant. <sup>a</sup> when compared with group I. <sup>b</sup> when compared with group II.

The activities of GSH, Vit C and Vit E level were significantly decreased in haemolysate of tumor bearing animals (Group II) as compared to control animals. Geraniol administrated rats showed significant increase in the non-enzymatic antioxidants levels (Group III and Group IV).

The activities of non-enzymatic antioxidants (GSH, Vit C and Vit E) level in liver of the experimental animals in each group are shown in Table 4.

The activities of SOD, CAT and GPx level significantly decreased in liver tissue of tumor bearing animals (Group II) as compared to control group. Geraniol administrated rats showed significant increase in the enzymatic antioxidants levels (Group III and Group IV). Geraniol administrated rats showed significant increase in the enzymatic antioxidants levels (Group III and Group IV).

**Table 2.** Effect of geraniol on the enzymatic antioxidants in liver of control and experimental animals

Parameters	Superoxide dismutase (SOD)	Catalase (CAT)	Glutathione peroxidase (GPx)
Group I (Control)	8.14 ± 0.47	60.88 ± 3.39	5.40 ± 0.30
Group II (DEN)	4.41 ± 0.34 <sup>a***</sup>	43.05 ± 1.56 <sup>a***</sup>	2.77 ± 0.42 <sup>a***</sup>
Group III (DEN + Geraniol)	7.30 ± 0.42 <sup>b***</sup>	54.23 ± 0.61 <sup>b***</sup>	4.43 ± 0.36 <sup>b***</sup>
Group IV (Geraniol)	7.87 ± 0.19 <sup>aNS</sup>	60.25 ± 1.15 <sup>aNS</sup>	5.16 ± 0.11 <sup>aNS</sup>

Values represent mean ± SD for rats in each group.

Units: SOD – units /mg protein

CAT – μ moles of H<sub>2</sub>O<sub>2</sub> utilized/min/mg protein.

GPx – μ moles of GSH oxidised /min/mg protein.

\*\*\*p<0.001; \*\* p<0.01; \* p<0.05

<sup>NS</sup> Non-significant. <sup>a</sup> when compared with group I. <sup>b</sup> when compared with group II.

**Table 3.** Effect of geraniol on the non-enzymatic antioxidants in haemolysate of control and experimental animals

Parameters	Glutathione reduced (GSH)	Vitamin C	Vitamin E
Group I (Control)	3.19 ± 0.29	1.90 ± 0.06	1.41 ± 0.11
Group II (DEN)	1.29 ± 0.04 <sup>a***</sup>	0.84 ± 0.03 <sup>a***</sup>	0.71 ± 0.04 <sup>a***</sup>
Group III (DEN+Geraniol)	2.76 ± 0.27 <sup>b***</sup>	1.79 ± 0.12 <sup>b***</sup>	1.26 ± 0.15 <sup>b***</sup>
Group IV (Geraniol)	2.85 ± 0.30 <sup>aNS</sup>	1.86 ± 0.04 <sup>aNS</sup>	1.24 ± 0.14 <sup>aNS</sup>

Values represent mean ± SD for 6 rats in each group.

Units: GSH, vitamin C, vitamin E- μg/mg protein.

\*\*\*p<0.001; \*\* p<0.01; \* p<0.05

<sup>NS</sup> Non-significant. <sup>a</sup> when compared with group I. <sup>b</sup> when compared with group II.

**Table 4.** Effect of geraniol on the non-enzymatic antioxidants in liver of control and experimental animals

Parameters	Glutathione reduced (GSH)	Vitamin C	Vitamin E
<b>Group I (Control)</b>	4.46 ± 0.35	2.81 ± 0.26	1.83 ± 0.05
<b>Group II (DEN)</b>	2.33 ± 0.16 <sup>a***</sup>	1.33 ± 0.08 <sup>a***</sup>	0.84 ± 0.02 <sup>a***</sup>
<b>Group III (DEN+Geraniol)</b>	3.83 ± 0.13 <sup>b***</sup>	2.45 ± 0.16 <sup>b***</sup>	1.72 ± 0.18 <sup>b***</sup>
<b>Group IV (Geraniol)</b>	4.20 ± 0.16 <sup>aNS</sup>	2.60 ± 0.30 <sup>aNS</sup>	1.80 ± 0.06 <sup>aNS</sup>

Values represent mean ± SD for 6 rats in each group.

Units: GSH, Vitamin C, Vitamin E- μg/mg protein

\*\*\*p<0.001; \*\* p<0.01; \* p<0.05

<sup>NS</sup> Non-significant. <sup>a</sup> when compared with group I. <sup>b</sup> when compared with group II.

## 4. Discussion

In the present study we demonstrate that the monoterpene geraniol has antioxidant activity. Geraniol, an acyclic monoterpene alcohol found in lemongrass and aromatic herb oils, has been shown to exert in vitro and in vivo antitumor activity against murine leukemia, hepatoma, and melanoma cells<sup>25-27</sup>.

M. S. Seema Farhath et al.<sup>28</sup> studied the Antioxidant activity of Geraniol, Geraniol acetate, Gingerol and Eugenol was employed by two complementary test systems, namely, 2,2-diphenyl-1-picrylhydrazyl (DPPH) free-radical scavenging and Super Oxide Dimutase (SOD) activity. Antioxidant activity of Gingerol was found to be higher than those of the others in DPPH assay.

Geraniol, a colourless liquid, is an acyclic terpene alcohol with a flowery rose-like odour. It is found widely as a chief constituent in essential oils including orange flower oil, lemon grass oil and lavender oil. Experimental studies demonstrated several pharmacological activities including antioxidant and anticancer potential of geraniol<sup>29</sup>.

Geraniol exerted anti-tumor activity against various cancer cells both in vitro and in vivo<sup>30-32</sup>. It has also been reported that geraniol exhibited potent insecticidal, antimicrobial and anti-inflammatory effects<sup>33,34</sup> reported that dietary geraniol suppressed hepatic HMG CoA reductase activity and lowered the levels of serum cholesterol in experimental animals.

Artega et al.<sup>35</sup> performed extend study focused on the antioxidant activity of low-molecular antioxidants present in the spice and some pharmaceuticals (gallic acid, sesamol, eugenol, thymol, carvacrol, vanillin, salicylaldehyde, limonene, geraniol, 4-hexylresorcinol).

Hanaa H et al.<sup>36</sup> studied the essential oil *Dracocephalan moldavica* L by TLC, GC-MS, HPLC and NMR and its constituents were geranyl acetate, geraniol, nerol, neryl acetate, neral and linolool. This oil showed antimicrobial and antioxidant activity.

S. Manoharan et al.<sup>37</sup> investigated the chemopreventive potential of geraniol, an acyclic monoterpene alcohol, by monitoring the tumor incidence and analyzing the status of phase II detoxification agents, lipid peroxidation by products and antioxidants in 7,12-dimethylbenz(a)anthracene (DMBA) induced mouse skin carcinogenesis and found that geraniol might have inhibited abnormal cell proliferation occurring in skin carcinogenesis by modulating the activities of phase II detoxification agents and through its free radical scavenging potential.

## 5. Conclusion

From this study Geraniol shows to exhibited chemopreventive potential against in N-Nitrosodiethylamine induced hepatocarcinogenesis in Wistar Albino Rats. Thus concluding that the chemopreventive potential of Geraniol could be due to its enzymatic antioxidant and non-enzymatic properties against N-Nitrosodiethylamine (DEN) induced Hepatocarcinogenesis (HCC).

## 6. References

1. Henricks PA, Nijkamp FP. Reactive oxygen species as mediators in asthma. *Pulm Pharmacol Ther.* 2001;14(6):409-20.
2. Singh B, Kakkar P. Mitochondria: a hub of redox activities and cellular distress control. *Mol Cell Biochem.* 2007; 305 (1-2):235-53.
3. Rahman I, Biswas SK, Kirkham PA. Regulation of inflammation and redox signaling by dietary polyphenols. *Biochem Pharmacol.* 2006; 72(11):1439-52.
4. Vandana S, Ram S, Ilavazhagan M, Kumar GD, Banerjee PK. Comparative cytoprotective activity of vitamin C, E and beta-carotene against chromium induced oxidative stress in marine macrophages. *Biomed. Pharmacother.* 2006; 60:71-6.
5. Karan M, Vasisht K, Handa SS. Antihepatotoxic activity of *Swertia chirata* on carbon tetrachloride-induced hepatotoxicity in rats. *Phytother. Res.* 1999; 13:24-30.
6. Chatterjee TK. Medicinal Plants with Hepatoprotective Properties. Herbal Options; Calcutta, India: Books and Applied Allied (P) Ltd; 2000.
7. Banu F. Antioxidant activity of geraniol against diethyl nitrosamine (den) induced hepatocarcinogenesis in adult wistar albino rats. *American J of Bio-pharm Biochem and Life Sci.* 2012 Mar; 1(Suppl 1):A37.
8. Mc-Intyre N, Rosalki S. Biochemical investigations in the management of liver disease. Prieto J, Rodes J, Shafritz, DA, editors. *Hepatobiliary Diseases.* Berlin: Springer- Verlag; 1992. p. 39-71.
9. Tricker AR, Pfundstein B, Theobald E, Preussman R, Spiengenthaler B. Mean daily intake of volatile N-Nitrosamines from foods and beverages in West Germany in 1989-90. *Food Chem Toxicol.* 1991; 29(11):729-32.
10. Craig WJ. Health-promoting properties of common herbs. *Am J Clin Nut.* 1999; 70(3):491S-99S.
11. Singh-Sangwan N, Sangwan RS, Luthra R, Thakur RS. Geraniol dehydrogenase: a determinant of essential oil quality in lemongrass. *Plant Med.* 2003; 59(2):168-70.
12. Sekiwa-Iijima Y, Aizawa Y, Kubota K. Geraniol dehydrogenase activity related to aroma formation in ginger (*Zingiber officinale*). *J Agric Food Chem.* 2001; 49(12):5902-06.

13. Iijima Y, Gang DR, Fridman E, Lewinsohn E, Pichersky E. Characterization of geraniol synthase from the peltate glands of sweet basil. *Plant Physiol.* 2004; 134(1):370–79.
14. Yang, T, Li J, Wang HX, Zeng Y. A geraniol-synthase gene from *Cinnamomum tenuipilum*. *Phytochem.* 2005; 66(3):285–93.
15. Ayoola GA, Johnson OO, Adelowotan T, Aibinu IE, Adenipekun E, Adepoju-Bellol AA, Coker HAB, Odugbemi TO. Evaluation of the chemical constituents and the antimicrobial activity of the volatile oil of *Citrus reticulata* fruit (Tangerine fruit peel) from South West Nigeria. *Afr J Biotechnol.* 2008; 7(13):2227–31.
16. F'guyer S, Afaq F, Mukhtar H. Photochemoprevention of skin cancer by botanical agents. *Photodermatol Photomed.* 2003; 19(2):56–72.
17. WHO FAS 52; 2004. Available from: [www.inchem.org/documents/jecfa/jecmono/v52je15.htm](http://www.inchem.org/documents/jecfa/jecmono/v52je15.htm); 2004.
18. Yu SG, Hildebrandt LA, Elson CE. Geraniol as inhibitor of mevalonate biosynthesis, suppresses the growth of hepatomas and melanomas transplanted to rats and mice. *J Nutr.* 1995; 125(11):2763–67.
19. Duncann RE, D Lau, El-Soheymy A, Archer MC. Geraniol and beta-ionone inhibit proliferation, cell cycle progression, and cyclin-dependent kinase 2 activity in MCF-7 breast cancer cells independent of effects on HMG-CoA reductase activity. *Biochem Pharmacol.* 2004; 68, 1739–47.
20. Carnesecchi S, Bras-Oncalves R, Bradaia A, Zeisel M, Gosse F, Poupon MF, Raul F. Geraniol, a component of plant essential oils, modulates DNA synthesis and potentiates 5-fluorouracil efficacy on human colon tumor xenografts. *Cancer Lett.* 2004; 215:53–9.
21. Liu YF, Zha BS, Zhang HL, Zhu XJ, Li YH, Zhu J, Guan XH, Feng ZQ, Zhang JP. Characteristic gene expression profiles in the progression from liver cirrhosis to carcinoma induced by diethylnitrosamine in a rat model. *J Exp Clin Canc Res.* 2009; 28:107. doi:10.1186/1756-9966-28-107
22. Tappel AL, Zalkin H. Inhibition of lipid peroxidation in mitochondria by vitamin E. *Arch Biochem Biophys* 1959; 80:333–336.
23. Marklund S, Marklund G. Involvement of superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem.* 1974; 47:469–74.
24. Dincer A, Aydemir T. Purification and characterization of catalase from chard. *J Enzyme Inhib.* 2001; 16(2):165–75.
25. Shoff SM, Grummer M, Yatvin MB, Elson CE. Concentration-dependent increase of murine P388 and B16 population doubling time by the acyclic monoterpene geraniol. *Cancer Res.* 1991; 51:37–42.
26. Yu SG, Hildebrandt LA, Elson CE. Geraniol, an inhibitor of mevalonate biosynthesis, suppresses the growth of hepatomas and melanomas transplanted to rats and mice. *J Nutr.* 1995; 125:2763–67.
27. Burke YD, Stark MJ, Roach SL, Sen SE, Crowell PL. Inhibition of pancreatic cancer growth by the dietary isoprenoids farnesol and geraniol. *Lipids.* 1997; 32:151–56.
28. Farhath MSS, Vijaya PP, Vimal M. Antioxidant activity of Geraniol, Geraniol acetate, Gingerol and Eugenol. *Research in Pharmacy.* 2013; 3(1):01–06.
29. Tiwari M, Kakkar P. Plant derived antioxidants - Geraniol and camphene protect rat alveolar macrophages against t-BHP induced oxidative stress. *Toxicol In vitro.* 2009; 23: 295–301.
30. Yu SG, Hildebrandt LA, Elson CE. Geraniol an inhibitor of mevalonate biosynthesis, suppresses the growth of hepatomas and melanomas transplanted to rats and mice. *J Nutr.* 1995; 125:2763–67.
31. Duncan RE, Lau D, El-Soheymy A, Archer MC. Geraniol and betaionone inhibit proliferation, cell cycle progression, and cyclindependent kinase 2 activity in MCF-7 breast cancer cells independent of effects on HMG-CoA reductase activity. *Biochem Pharmacol.* 2004; 68:1739–47.
32. Carnesecchi S, Bras-Goncalves R, Bradaia A, Zeisel M, Gosse F, Poupon MF, Raul F. Geraniol, a component of plant essential oils, modulates DNA synthesis and potentiates 5-fluorouracil efficacy on human colon tumor xenografts. *Cancer Lett.* 2004; 215(1):53–59.
33. Chen W, Viljoen AM. Geraniol - A review of a commercially important fragrance material. *S Afr J Bot.* 2010; 76(4):643–51.
34. Crowell PL. Prevention and therapy of cancer by dietary monoterpenes. *J Nutr.* 1999; 129:775–78.
35. Arifullah M, Namsa ND, Gopa GR et al. Evaluation of anti-bacterial and anti-oxidant potential of andrographolide and echiodinin isolated from callus culture of *Andrographis paniculata* Nees. *Asian Pacific Journal of Tropical Biomedicine.* 2013 Aug; 3(8):604–10.
36. El-Baky HHA, El-Baroty GS. Chemical and biological evaluation of the essential oil of egyptian moldavian balm. *Global J Biotechnol Biochem Res.* 2007; 2(2):74–80.
37. Manoharan S, Selvan MV. Chemopreventive potential of geraniol in 7,12-dimethylbenz(a) anthracene (DMBA) induced skin carcinogenesis in Swiss albino mice. *J Environ Biol.* 2012; 33:255–60.